


## Problem set 3

### 3.1 The Aharonov-Bohm effect

We consider a double slit experiment as sketched in the figure. Electrons are emitted from a source  $S$  and can pass through one of the two slits of a first screen before being registered on a second screen. When a large number of electrons are registered they are found to form an interference pattern with minima and maxima on the screen.



Behind the middle part of the first screen a solenoid is placed which carries a magnetic flux  $\Phi$ . The direction of the solenoid is parallel to the direction of the two slits, so that the paths through the upper slit pass on one side of the solenoid and the paths through the lower slit pass on the other side of the solenoid. We consider the magnetic field to be completely screened from the region where the electrons move, so that at no point along the trajectories of the electron there is a magnetic force acting on the particles. Nevertheless, quantum theory predicts that the strength of the magnetic flux will influence the interference pattern so that the maxima and minima are shifted up or down when the flux is changed. This is called the Aharonov-Bohm effect.

We consider in the following the distance  $d$  between the screens and the distance  $D$  between the source and the first screen to be much larger than the distance  $a$  between the two slits, and also to be much larger than the distance  $y$  from the central point of the second screen to any point  $P$  where an electron is registered.

As a reminder the classical Lagrangian of an electron moving in a magnetic field is

$$L(\mathbf{r}, \dot{\mathbf{r}}) = \frac{1}{2}m\dot{\mathbf{r}}^2 + e\mathbf{A}(\mathbf{r}) \cdot \dot{\mathbf{r}} \quad (1)$$

with  $\mathbf{A}$  as the vector potential, and the magnetic field thus given as  $\mathbf{B} = \nabla \times \mathbf{A}$ . As follows from Stokes' theorem the magnetic flux is given as the line integral

$$\Phi = \oint_C \mathbf{A} \cdot d\mathbf{r} \quad (2)$$

where  $C$  is any given closed loop that encircles once the solenoid.

We consider the situation where a single electron is emitted at time  $t = 0$  from the source and is registered at a later time  $t$  at a point  $P$  of the screen. The probability distribution over the screen for

where the electron is registered can be written as

$$p(y) = \lambda |\mathcal{G}(\mathbf{r}_P, t; \mathbf{r}_S, 0)|^2 \quad (3)$$

with  $y$  as the vertical coordinate of P,  $\lambda$  as a proportionality factor and  $\mathcal{G}(\mathbf{r}_P, t; \mathbf{r}_S, 0)$  as the propagator from the initial point  $(\mathbf{r}_S, 0)$  to the final point  $(\mathbf{r}_P, t)$ .

We consider in the following the semi-classical approximation to the propagator, which we write as

$$\mathcal{G}(\mathbf{r}_P, t; \mathbf{r}_S, 0) = N \sum_{n=1}^2 e^{\frac{i}{\hbar} S_n} \quad (4)$$

where  $S_n$  is the action integral for classical free-particle motion either through the upper slit ( $n = 1$ ) or through the lower slit ( $n = 2$ ), and  $N$  is a ( $y$ -dependent) normalization factor which is assumed to be independent of the path. Since the classical motion is not affected by the magnetic field, both  $\lambda$  and  $N$  are independent of the magnetic flux.

a) Show that the probability  $p(y)$  depends on the *difference* between the action integrals of the two paths.

b) Show that the difference between the two action integrals can be written as a function of the magnetic flux  $\Phi$ .

c) Show that the probability  $p(y)$  depends periodically on the magnetic flux  $\Phi$ . What is the flux period? Describe qualitatively how the interference pattern changes with variations in  $\Phi$ .

### 3.2 Spin operators and Pauli matrices

A spin half operator  $\hat{\mathbf{S}}$  is defined in the standard way as

$$\hat{\mathbf{S}} = \frac{\hbar}{2} \boldsymbol{\sigma} \quad (5)$$

where  $\boldsymbol{\sigma}$  is a vector with the three Pauli matrices  $(\sigma_1, \sigma_2, \sigma_3)$  (or equivalently written as  $(\sigma_x, \sigma_y, \sigma_z)$ ) as Cartesian components. We use the standard expressions for these 2x2 matrices, as given in the lecture notes. We also introduce the rotated Pauli matrix, defined by  $\sigma_{\mathbf{n}} = \mathbf{n} \cdot \boldsymbol{\sigma}$ , where  $\mathbf{n}$  is an unspecified three dimensional unit vector.

a) Show that  $\sigma_{\mathbf{n}}$  has eigenvalues  $\pm 1$ , and the eigenstate (in matrix form) corresponding to the eigenvalue  $+1$  is (up to an arbitrary phase factor)

$$\Psi_{\mathbf{n}} = \begin{pmatrix} \cos \frac{\theta}{2} \\ e^{i\phi} \sin \frac{\theta}{2} \end{pmatrix} \quad (6)$$

with  $(\theta, \phi)$  as the polar angles of the unit vector  $\mathbf{n}$ . Also show the relation

$$\Psi_{\mathbf{n}}^\dagger \boldsymbol{\sigma} \Psi_{\mathbf{n}} = \mathbf{n} \quad (7)$$

b) Show, by using operator identities from Problem Set 1, the following relation

$$e^{-\frac{i}{2}\alpha\sigma_z} \sigma_x e^{\frac{i}{2}\alpha\sigma_z} = \cos \alpha \sigma_x + \sin \alpha \sigma_y \quad (8)$$

Explain why this shows that the unitary matrix

$$\hat{U} = e^{-\frac{i}{2}\alpha\sigma_n} = e^{-\frac{i}{\hbar}\alpha\mathbf{n}\cdot\hat{\mathbf{S}}} \quad (9)$$


induces a spin rotation of angle  $\alpha$  about the axis  $\mathbf{n}$ .

c) Demonstrate, by expansion of the exponential function, the following identity

$$e^{-\frac{i}{2}\alpha\sigma_n} = \cos \frac{\alpha}{2} \mathbb{1} - i \sin \frac{\alpha}{2} \sigma_n \quad (10)$$

with  $\mathbb{1}$  as the 2x2 identity matrix.

### 3.3 Oscillations in ammonia molecules (Midterm Exam 2010)



The ammonia molecule has the chemical formula  $NH_3$ , which means that it is composed of three hydrogen and one nitrogen atoms. The left part of the figure shows the spatial structure of the molecule, where the hydrogen atoms define a planar, equilateral triangle (in the  $yz$ -plane) and the nitrogen atom is located on the orthogonal symmetry axis ( $x$ -axis) at some distance from the plane of the hydrogen atoms.

With the plane of the hydrogen atoms being fixed, there are, however, two possible positions of the nitrogen atom that are equally favored with respect to potential energy. They are located symmetrically about the plane, as indicated in the figure.

In the quantum description we associate two different state vectors  $|\psi_R\rangle$  and  $|\psi_L\rangle$  with these two positions of the nitrogen atom. In the right part of the figure the situation is pictured with the potential energy and the two wave functions shown as functions of the position of the nitrogen atom along the symmetry axis. The potential has the form of a double well with two degenerate ground state positions. With  $\hat{H}$  as the Hamiltonian we write this degeneracy as

$$\langle\psi_L|\hat{H}|\psi_L\rangle = \langle\psi_R|\hat{H}|\psi_R\rangle \equiv E_0 \quad (11)$$

There is however a correction to this picture. Even though there is potential barrier between the two equilibrium positions, there is a small probability for quantum tunneling from one position to the other. This is represented by a non-vanishing matrix element

$$\langle \psi_L | \hat{H} | \psi_R \rangle \equiv \lambda \quad (12)$$

were we may assume  $\lambda$  to be real and positive. The value of this matrix element is very small, which means that the corresponding transition time from one minimum of the potential to the other is very long, but the result is that if the nitrogen atom initially is in one of the wells it will oscillate back and forth between the two minima at a low frequency (compared to other atomic frequencies).

The true ground state is however a stationary state, which to a good approximation is a superposition of the states  $|\psi_L\rangle$  and  $|\psi_R\rangle$  associated with the two minima. In the following we restrict the description to the two-dimensional Hilbert space spanned by these two vectors.

a) Write the Hamiltonian as a  $2 \times 2$  matrix and find the energy eigenvalues  $E_0^\pm$  and eigenstates  $|\psi_0^\pm\rangle$ , when the  $\lambda$  terms are included. Express the ground state  $|\psi_0^-\rangle$  and the excited state  $|\psi_0^+\rangle$  as linear combinations of  $|\psi_L\rangle$  and  $|\psi_R\rangle$  and describe briefly with words the characteristics of the two energy eigenstates.

The ammonia molecule has an electric dipole moment which arises from the tendency of the nitrogen atom to attract an electron from the hydrogen atoms. The dipole moment is directed along the symmetry axis in the opposite direction of the nitrogen atom. We assume now that the ammonia molecule is located in a constant electric field  $\mathcal{E}$  directed along the x-axis. The field introduces a new term  $\hat{H}_d$  in the Hamiltonian with matrix elements

$$\begin{aligned} \langle \psi_L | \hat{H}_d | \psi_L \rangle &= -\langle \psi_R | \hat{H}_d | \psi_R \rangle \equiv \Delta \\ \langle \psi_L | \hat{H}_d | \psi_R \rangle &= \langle \psi_R | \hat{H}_d | \psi_L \rangle = 0 \end{aligned} \quad (13)$$

with  $\Delta = \mathcal{E} d$ , and with  $d$  as the electric dipole moment of the molecule.

b) Determine the new energy eigenvalues  $E_\pm$  with this additional term in the Hamiltonian, and make a plot that shows how the two energy levels change with variable  $\Delta$  from a large negative to a large positive value (from  $\Delta \ll -\lambda$  to  $\Delta \gg \lambda$ ). Choose  $E_\pm/\lambda$  and  $\Delta/\lambda$  as variables.

c) Determine eigenvectors  $|\psi_\pm\rangle$  expressed in terms of  $|\psi_L\rangle$  and  $|\psi_R\rangle$  and plot, as functions of  $\Delta$ , the overlaps  $|\langle \psi_L | \psi_\pm \rangle|^2$  between the energy eigenvectors and  $|\psi_L\rangle$ .

The situation we have here is sometimes referred to as an *avoided crossing* between the two energy levels. Give a brief qualitative description of the crossing based on the plotted curves.

We next assume the electric field to vary periodically with time,  $\mathcal{E} = \mathcal{E}_0 \cos \omega t$ , and correspondingly,  $\Delta = \Delta_0 \cos \omega t$ , with  $\Delta_0$  as a positive constant.

d) Show that in the  $\{|\psi_0^\pm\rangle\}$  basis the Hamiltonian can be expressed as

$$\hat{H} = E_0 \mathbb{1} + \lambda \sigma_z + \Delta_0 \cos \omega t \sigma_x \quad (14)$$

with  $\sigma_x$  and  $\sigma_z$  as standard Pauli matrices.

The last term in (14) can be written as

$$\Delta_0 \cos \omega t \sigma_x = \frac{1}{2} \Delta_0 (e^{i\omega t} \sigma_- + e^{-i\omega t} \sigma_+) + \frac{1}{2} \Delta_0 (e^{-i\omega t} \sigma_- + e^{+i\omega t} \sigma_+) \quad (15)$$

where  $\sigma_{\pm} = \frac{1}{2}(\sigma_x \pm i\sigma_y)$  can be viewed as raising and lowering operators in the spectrum of the two-level Hamiltonian. Assuming  $\omega$  to be positive the first term will usually give the most important contribution to the Hamiltonian. This motivates the so-called *rotating wave approximation*, where the last term in (15) is omitted. In the following apply this approximation with the Hamiltonian given by

$$\hat{H} = E_0 \mathbb{1} + \lambda \sigma_z + \frac{1}{2} \Delta_0 (e^{i\omega t} \sigma_- + e^{-i\omega t} \sigma_+) \quad (16)$$

e) Show that this has the same form as the spin Hamiltonian in a rotating magnetic field, discussed in Sect.1.3.2 of the lecture notes. Outline the method used to find the time evolution operator and give the expressions for the Rabi frequency  $\Omega$  and resonance frequency  $\omega_0$  in terms of the parameters  $\lambda$  and  $\Delta_0$ . It may be convenient here to re-define the zero-point of the energy so that  $E_0 = 0$ . Comment on why the value of  $E_0$  is not important. It is sufficient to refer to results from the lecture notes without a detailed derivation.

f) Initially, at time  $t = 0$ , the system is in the left shifted state  $|\psi_L\rangle$ . Determine the time dependence of the overlap of the time evolved state  $|\psi(t)\rangle$  with the right shifted state,  $\langle\psi_R|\psi(t)\rangle$ .

g) Assuming that the strength of the oscillating field is given by  $\Delta_0 = 2\lambda$ , examine numerically the time dependent function  $|\langle\psi_R|\psi(t)\rangle|^2$  by making a plot over several periods of this function, for two different values of the frequency, 1) at resonance,  $\omega = \omega_0$  and 2) off resonance with  $\omega = \omega_0/10$ . Use the dimensional variable  $\tau = 2\pi\lambda t$  as time coordinate. Make a (qualitative) discussion of what the curves show and compare with the related curve in the case when the electric field is turned off,  $\Delta_0 = 0$ .