

AST5770

Solar and stellar physics

Sven Wedemeyer, University of Oslo, 2025

Solar filaments, prominences, CMEs Recap

Filaments and prominences — recap

Filament on disk — prominence at the limb


1. What is the difference between a filament and a prominence?
2. What are the main three types?
3. How long can filaments be?
4. Why are filaments dark and prominences bright?
5. How can cool dense plasma “just hang up there”?
6. What is the plasma temperature in filaments and how does that compare to their environment?
7. Can you explain the following parts of filaments?
 1. Leg
 2. Spine
 3. Barb
8. Why do erupting filaments often show an unwinding motion?
9. How long does it take for ejected plasma to reach Earth?

Magnetic field in the solar atmosphere

Recap — Magnetism

Magnetic pressure

- Ionised gas (plasma) in motion — electric and magnetic fields need to be considered
- Additional contributions from magnetic pressure inside magnetic flux concentration
- **Magnetic pressure** $P_m = B^2 / 8\pi$
- Pressure balance $\Rightarrow$ lower gas pressure inside the flux concentration than outside
- Gas pressure of surrounding drops with height $\Rightarrow$ Magnetic structure funnels out (wine-glass shape)



- $B_2 > B_1$

- $P_2 < P_1$

- $\rho_2 < \rho_1$


Recap — Magnetism


Plasma-Beta

- Plasma- β describes the ratio of thermal to magnetic pressure

$$\beta = \frac{P_g}{P_m} = \frac{8\pi P_g}{B^2}$$

- $\beta < 1$: Magnetic field dominates** and dictates the dynamics of the gas
- $\beta > 1$: Thermal gas dynamics dominate** and forces the field to follow — The magnetic field is **frozen-in**.
- β is a local quantity but the typical range of values changes with radius:
 - Convection zone: $\beta > 1$
 - Lower atmosphere (outside strong magnetic field concentrations): $\beta > 1$
 - Chromosphere: transition to $\beta < 1$
 - Corona: $\beta \ll 1$







Magnetic features in the solar atmosphere

(smaller than sunspots)

The solar atmosphere outside from Active Regions



Credit to Big Bear Solar Observatory and NASA for data and video content.
Credit to Tanmoy Samanta and Hui Tian from Peking University for video production.



← 10 000 km →

granulation

Photosphere of the Sun = $\tau(\lambda=500\text{nm}) = 1$
narrow layer where visible continua become optically thick

G-band (430nm)

“Quiet Sun”

- Away from active regions with strong magnetic fields
- Granulation pattern (as result of thermal convection at the solar “surface”)
- Magnetic elements

Sunspot

Umbra


Penumbra

Magnetic pores



Magnetic field in the solar atmosphere

Pores

- Umbra only — like sunspots but without a penumbra.
- Appear when a strong magnetic field emerges through solar surface (inhibits convective transfer of heat from below as in sunspots, appears dark)
- “Small” (1000 - 6000 km in diameter) but can include smaller bright structures (convective heat transfer from below not completely suppressed there)
- Dispersed magnetic fields around pores can produce bright points and chains in intergranular lanes in surrounding





Solar pores
VTT 2012
G-band 430 nm.



Magnetic field in the solar atmosphere

Photospheric magnetic field


Magnetograms (via Stokes V at 632nm from Spectro-Polarimeter (SP) onboard Hinode/SOT



Magnetic field in the solar atmosphere

Magnetic field as function of spatial scale


- (Average) magnetic field strength measured for structures ("flux tubes") with different sizes
 - From large sunspots to the smallest detectable magnetic elements
- Surprisingly constant field strength per area
- Remember: Magnetic fields "compete" with thermal pressure (and convective motions)
 - ➔ Equipartition field strength sets limits



Magnetic field in the solar atmosphere



Magnetic field as function of spatial scale

- Observable consequences strongly depend on area covered by magnetic field structure
 - Large: darker than surrounding
 - Small: brighter (!) than surrounding
- Remember: Sunspots — strong magnetic fields impede convective heat transport from below, resulting in lower temperature
- **What about small magnetic elements?**



Magnetic field in the solar atmosphere

Photospheric magnetic field in Quiet Sun regions





- **Remember:**

1. Magnetic field **emerges** through the "surface" (photosphere)
 - Magnetic field structure on a large range of spatial scales
2. Field is **advected**

Magnetic field in the solar atmosphere

Photospheric magnetic field in Quiet Sun regions




- **Remember:**

1. Magnetic field **emerges** through the "surface" (photosphere)
 - Magnetic field structure on a large range of spatial scales
2. Field is **advected**

Magnetic field in the solar atmosphere

Advection — supergranulation scales


- Magnetic field emerges to the surface
 - Away from strong fields (sunspots):
High plasma- β in the photosphere
 - **Frozen-in** magnetic field
 - Field is advected with the photospheric velocity field towards the **edges of supergranules**
 - Concentrated there, resulting in stronger magnetic flux concentrations
 - Observable as **magnetic network**
 - Encloses **inter-network regions**
- Magnetogram (grayscale)
 - Horizontal flow field (arrows)
 - Supergranule boundaries: yellow



Magnetic field in the solar atmosphere

Advection — magnetic field on granulation scales

- Advection into intergranular lanes (downflow lanes between granules)
- Concentration into stronger flux concentrations but fewer than in the network




Granulation image,
Fe I 630.25 nm line

Overlaid
magnetogram
contours

30, 50, 70 and 90 G

(Dominguez Cerdena
et al., 2003)



Magnetic field in the solar atmosphere

Photospheric magnetic field in Quiet Sun regions

- Distribution of longitudinal magnetic fluxes observed with different instruments
(Hinode/NFI magnetograms, SOHO/MDI high resolution + full disk magnetograms)
 - Magnetic flux value vs. its occurrence scales follows a power law (=linear in log-log)
- ➡ Scalable phenomena
- ➡ Beyond current detection limit?


Magnetic flux Φ

(Flux = per area!)

cgs unit: Maxwell (Mx)


1 Mx = 1 G cm⁻²

Longitudinal = component parallel (along) magnetic field vector





Magnetic field in the solar atmosphere

Photospheric magnetic field in internetwork regions




- Average field strength = 170 G, FWHM of distribution = 190 G.

- Flux tubes with $B > 1$ kG occur but not dominant
- More of those in the magnetic network



- Magnetic field inclination: highly inclined fields very abundant!
 - Maximum at 90° (horizontal fields)
 - Bumps near 0° and 180° : vertical fields in intergranular lanes


Magnetic field in the solar atmosphere



G-band observation — magnetic field concentrations visible with high contrast in this band

Mag. field




Flux emergence



Magnetic field in the solar atmosphere

Flux emergence

Top
View



Side
View




One Million
Kelvin



Ten Million
Kelvin


Magnetic field in the solar atmosphere

Flux emergence




- Emergence of bipolar regions with a large range of contained magnetic flux
 - Many regions with little flux, fewer with a lot of flux
- Varies over solar cycle

Ephemeral regions = short-lived, small bipolar regions (do not develop sunspots)




Magnetic field in the solar atmosphere

Flux emergence



- Emergence of bipolar regions with a large range of contained magnetic flux
- Many regions with little flux, fewer with a lot of flux
- Varies over solar cycle


Ephemeral regions = short-lived, small bipolar regions (do not develop sunspots)



Magnetic field in the solar atmosphere


Structure of Quiet Sun regions

Magnetic field extrapolation from photospheric magnetograms



Jafarzadeh et al 2021

Observed: Fibrils in Ca II K




Peter et al 2021

Magnetic field in the solar atmosphere

Numerical simulation

- Small part of an internetwork region




The solar atmosphere

Radiative transfer effects

Radiative transfer effects


- Remember: Magnetic pressure counterbalances thermal (gas) pressure
- $B_i \gg B_o$
- ➔ Lower thermal pressure inside region with strong magnetic field $P_{g,i} \ll P_{g,o}$
- ➔ Lower gas density (=fewer atoms) $\rho_i \ll \rho_o$
- ➔ **Lower opacity** inside the magnetic flux structure
- Remember: Optical depth according to opacity along line of sight



- ➔ **Optical depth lower** inside magnetic field structure than outside
- ➔ Looking deeper into the Sun inside the magnetic field structure
- Lower height of optical depth unity = **Wilson depression** (in sunspots),

Radiative transfer effects

- Effect also visible for weaker magnetic field structures on smaller spatial scales
- Difference: Convective energy transport not suppressed (as much) as under sunspots!




Radiative transfer effects

- Observing at inclination angle $\theta > 0$ (oblique)
($\mu = \cos \theta$)
- ➡ Geometrically: Longer line of sight path element per change in height
- ➡ “Picking up” more opacity already higher up in atmosphere due to projected

Consequence 1 (outside magnetic field structures)

- Optical depth unity reached already higher up!
- View directly from top allows look deepest into atmosphere
- **Mapped height** in the atmosphere (i.e. z where $\tau=1$) **increases with viewing angle θ**




Radiative transfer effects

- Observing at inclination angle $\theta > 0$ (oblique)
($\mu = \cos \theta$)
- ➡ Geometrically: Longer line of sight path element per change in height
- ➡ “Picking up” more opacity already higher up in atmosphere due to projected
- ➡ Now looking through a “thin flux tube”

Consequence 2:

- Views deeper into the atmosphere through the flux tube
- Temperature is higher deeper down
- Higher temperature sampled at inclined viewing angle but only on the side where one sees through the flux tube



➡ **Hot wall effect:** Magnetic feature appears to be brighter on one side!


Radiative transfer effects

- View from top $\theta = 0$
- ➡ “Picking up” less opacity when looking through a “thin flux tube” from top

Consequence 3:

- Views deeper into the atmosphere through the flux tube
- Temperature is higher deeper down
- Higher temperature sampled
- Magnetic feature appears bright!



➡ **Magnetic bright point**



Radiative transfer effects

Inclination angle

$$\mu = \cos \theta$$




Radiative transfer effects

Limb darkening

In the **visible continuum**:


- Solar disk darker at the edge (limb) than at the centre of the solar disk!
- Decreasing brightness implies decreasing temperature (assuming simple blackbody radiation)
- Remember: Observation at increasing angle θ (=decreasing μ) maps layer at increasing height in the atmosphere
- ➔ Temperature decreases with height for the height range mapped by this continuum range!
- Note: Optical depth unity reached at bottom of photosphere (basically defines it) in visible continuum, "looks" deepest into the atmosphere



Radiative transfer effects

Limb brightening



- In the **mm continuum**:
Solar disk **brighter** at the edge (limb)
than at the centre of the solar disk
(on average, do not look at ARs)!
- **Increasing** brightness implies
increasing temperature (assuming
simple blackbody radiation)
- Remember: Observation at increasing
angle θ (=decreasing μ) maps layer at
increasing height in the atmosphere
➔ Temperature **increases** with height
for the height range mapped by this
continuum range!
- Note: Optical depth unity reached in
the chromosphere, i.e. much higher
than the visible continuum



Radiative transfer effects

Centre-to-limb variation (CLV)


- Centre-to-limb variation depends on mapped height range in the atmosphere
 - Photosphere: darkening
 - Chromosphere: brightening
 - Opposite gradient of the temperature stratification!
- CLV measured from observations help to construct (1D) average atmosphere models (even for different types of region from Quiet Sun to sunspot umbra)



Radiative transfer effects

Centre-to-limb variation (CLV)


- 3D numerical simulations for different stellar types
- Intensity maps for as function of inclination angle μ from disk-center to limb (visible continuum).
- All show limb darkening.
- Observational imprint of photospheric temperature stratification!



Radiative transfer effects


Faculae

- Faculae = **bright** areas most easily seen near solar limb (hot wall effect!)
- Areas with (small-scale) magnetic elements (in photosphere)
- Area on Sun covered with faculae varies over solar cycle
- Corresponding increase in brightness (across whole disk) more than compensates for darkening by sunspots!
- $\sim 0.1\%$ brighter at solar maximum than at solar minimum
- **Plage** = chromospheric counterpart of a facular region



The solar atmosphere

Different regions — photosphere


(Average) magnetic field strength	Active Region (Large) area with strong magnetic field	Sunspot Areas of concentrated very strong field, appear dark	Umbra Central compact part, dark Penumbra Surrounding, filamentary
		Faculae bright (filamentary) areas	
	Quiet Sun Outside Active Regions, weaker magnetic field	Network Concentrations of strong magnetic field, filamentary/mesh-like	
		Inter-network Areas with weak magnetic field inside network cells	
			

The solar atmosphere

Different regions — chromosphere


(Average) magnetic field strength

Active Region (Large) area with strong magnetic field	Sunspot Areas of concentrated very strong field, appear dark	Umbra Central compact part, dark Penumbra Surrounding, filamentary
	Plage bright area, higher temperature, often proceeds formation of sunspots	
Quiet Sun Outside Active Regions, weaker magnetic field	Network Concentrations of strong magnetic field, filamentary/ mesh-like	
	Inter-network Areas with weak magnetic field inside network cells	





The solar atmosphere



- Different parts of the line formed at different heights
- Looking a bit higher in the atmosphere
- Spatial scales corresponding to granulation visible
- Prominent scale with super granulation, here with cell sizes of $\sim 30\text{Mm}$
- Extension of magnetic field from photospheric footprints into the chromosphere



The solar atmosphere





The solar atmosphere



Magnetic field in the solar atmosphere


Structure of Quiet Sun regions



Magnetic field in the solar atmosphere

Structure of Quiet Sun regions


- Modern telescopes with high spatial + temporal + spectral show a new picture of the "Quiet" Sun
- Dynamic intermittent structure across many scales, plethora of physical processes



Magnetic field in the solar atmosphere

Structure of Quiet Sun regions

- Modern telescopes with high spatial + temporal + spectral show a new picture of the "Quiet" Sun
- Dynamic intermittent structure across many scales, plethora of physical processes



The solar atmosphere

Structure of Quiet Sun regions

- Magnetic field in the photosphere: Footpoints with vertical field
- Chromosphere: Magnetic field connects polarities, forms loops (horizontal field, "canopy")
- Smaller loops can connect lower (small-scale canopies, horizontal field in photosphere)
- Different diagnostics (spectral lines/continua) show different layers and aspects
 - Horizontal chromospheric field clearer at some wavelengths (e.g. : H α core) than at others

