University of Oslo

FYS4170/9170 — Relativistic Quantum Field Theory

Problem set 8

Problem 1 Wick's theorem and contractions (J. Skaar)

When computing correlation functions, it is useful to decompose the fields into positive and negative frequency parts. For the field in the interaction picture, this amounts to

$$\phi(x) = \phi^+(x) + \phi^-(x) \tag{1}$$

with

$$\phi^+(x) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\mathbf{p}}}} a_{\mathbf{p}} e^{-ipx} \tag{2}$$

$$\phi^-(x) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\mathbf{p}}}} a^\dagger_{\mathbf{p}} e^{ipx} \tag{3}$$

such that

$$\phi^+(x)|0\rangle = 0\,, \qquad \langle 0|\phi^-(x) = 0\,. \tag{4}$$

a) Prove Wick's theorem for two fields:

$$T\phi(x_1)\phi(x_2) = N\{\phi(x_1)\phi(x_2) + \overbrace{\phi(x_1)\phi}(x_2)\}, \tag{5}$$

where

$$\overbrace{\phi(x_1)\phi}(x_2) = \begin{cases} [\phi^+(x_1), \phi^-(x_2)], & \text{if } x_1^0 > x_2^0, \\ [\phi^+(x_2), \phi^-(x_1)], & \text{if } x_2^0 > x_1^0. \end{cases} \tag{6}$$

Also prove that

$$\overbrace{\phi(x_1)\phi}(x_2) = \langle 0|T\phi(x_1)\phi(x_2)|0\rangle = D_F(x_1 - x_2). \tag{7}$$

b) Use Wick's theorem to obtain an expression for the vacuum expectation value

$$\langle 0|T\{\phi(x_1)\phi(x_2)\cdots\phi(x_m)\}|0\rangle. \tag{8}$$


What is the value of the vacuum expectation value if there is an odd number m of fields?

c) Calculate $\langle\Omega|\phi(x)|\Omega\rangle$ for ϕ^4-theory. Here $\phi(x)$ is the field in the Heisenberg picture (not interaction picture as above!).


d) Go through the induction proof of Wick's theorem in P&S, p. 90. Be aware of the "paradox" $1 = N\{1\} = N\{aa^\dagger - a^\dagger a\} = a^\dagger a - a^\dagger a = 0$, which tells us that it is not allowed to insert the value of a commutator inside the $N\{\cdot\}$.

Problem 2 Feynman diagrams and symmetry factors (T. Klungland)

This problem explores some fully connected diagrams[1] contributing to the ϕ^4 4-point function, denoted by $\langle\Omega|T\{\phi(x_1)\phi(x_2)\phi(x_3)\phi(x_4)\}|\Omega\rangle$. At the leading order, it is given by the diagram[2]



At order $\mathcal{O}(\lambda^2)$ there are three contributing diagrams:



We denote these by S, T, and U respectively.

a) Using the Feynman rules on page 94 in P&S, write down the position space representation of each of these diagrams. Do *not* attempt to perform the integrals (unless you have entirely too much spare time), simply leave them undetermined.

b) The symmetry factors of the diagrams in the previous part can be derived by Taylor expanding the numerator of Eq. (4.31) in P&S (we have four external fields instead of two; otherwise the equation can be used directly). At $\mathcal{O}(\lambda^2)$ we have, dropping the I subscripts,

$$\frac{1}{2!}\left(\frac{-i\lambda}{4!}\right)^2 \int d^4z d^4w \langle 0|T\big\{\phi(x_1)\phi(x_2)\phi(x_3)\phi(x_4)\phi^4(z)\phi^4(w)\big\}|0\rangle,$$


where the factor 1/2! appears due to the Taylor expansion. The symmetry factor is found by counting the number of equivalent contractions corresponding to each particular diagram; when multiplying the above expression by this number, the symmetry factor s can be read off from the prefactor to the integral as $-\lambda^2/s$.

Use this to explicitly verify the symmetry factor of the S diagram (the process for the other two is identical).

[1] In a *fully connected* diagram all lines are connected to every external point by a sequence of lines and vertices. As you will see later in the course, these are the only diagrams that end up contributing to the correlation functions, as all disconnected diagrams factor out in both the numerator and denominator of Eq. (4.31) in P&S and thus cancel.

[2] In all diagrams in this exercise we place dots on the vertices in order to make it more explicit when two lines "accidentally" cross each other and when there is a genuine vertex.

c) Repeat the above process for a sample $\mathcal{O}(\lambda^3)$ diagram, given by



i.e. find the symmetry factor of the diagram both directly by looking at the symmetries of the diagram, and from counting contractions.