

IN3070/4070 – Logic – Autumn 2019

Lecture 5: Soundness & Completeness for 1st-order LK

Martin Giese

18th September 2019



DEPARTMENT OF
INFORMATICS



UNIVERSITY OF
OSLO

Today's Plan

- ▶ Preliminaries and Reminders
- ▶ Soundness Proof
- ▶ Completeness: Preliminaries
- ▶ Proof of Completeness
- ▶ Examples of Counter-model Construction

Outline

- ▶ Preliminaries and Reminders
- ▶ Soundness Proof
- ▶ Completeness: Preliminaries
- ▶ Proof of Completeness
- ▶ Examples of Counter-model Construction

Reminder Soundness of LK

- ▶ We want all LK-provable sequents to be valid!
- ▶ If they are not, then LK would be **incorrect** or **unsound** ...

Definition 1.1 (Soundness).

*The sequent calculus LK is **sound** if every LK-provable sequent is valid.*

Theorem 1.1.

The sequent calculus LK is sound.

Assumptions about the first order language

- ▶ We assume that a first-order language is given, by sets of constants, function symbols, and predicates.
- ▶ Some rules require “fresh” constants, so we assume that the set of constant symbols \mathcal{A} is (countably) infinite.
- ▶ A root sequent $\Gamma \Longrightarrow \Delta$ consists of *closed* formulae.
- ▶ We show that if $\Gamma \Longrightarrow \Delta$ is provable, then $\Gamma \Longrightarrow \Delta$ is valid

Reminder: Semantics for Sequents

Definition 1.2 (Valid sequent).

A sequent $\Gamma \Longrightarrow \Delta$ is *valid* if all interpretations that satisfy all formulae in Γ satisfy at least one formula in Δ .

Definition 1.3 (Countermodel/falsifiable sequent).

- ▶ An interpretation \mathcal{I} is a *countermodel* for the sequent $\Gamma \Longrightarrow \Delta$ if $v_{\mathcal{I}}(A) = T$ for all formulae $A \in \Gamma$ and $v_{\mathcal{I}}(B) = F$ for all formulae $B \in \Delta$
- ▶ We say that a countermodel for a sequent *falsifies* the sequent.
- ▶ A sequent is *falsifiable* if it has a countermodel.

Syntax vs. Semantics for Quantifiers

- ▶ Soundness and Completeness give the connection between
 - ▶ syntax (= calculus)
 - ▶ semantics ($\mathcal{I} \models \varphi$)
- ▶ Quantifier rules use substitutions
- ▶ The semantics of quantifiers use variable assignments
- ▶ We therefore need a connection between
 - ▶ substitutions (= syntactic operations)
 - ▶ variable assignments (= semantic objects)
- ▶ This connection is given by the substitution lemma

Reminder: Substitution Lemma

Theorem 1.2 (Substitution Lemma for Formulae).

Given an interpretation $\mathcal{I} = (D, \iota)$ and a variable assignment α for \mathcal{I} . Given also a variable $y \in \mathcal{V}$, a formula A and a term $s \in \mathcal{T}$, such that $\{y \setminus s\}$ is capture-free for A .

$$v_{\mathcal{I}}(\alpha, A[y \setminus s]) = v_{\mathcal{I}}(\alpha \{y \leftarrow v_{\mathcal{I}}(\alpha, s)\}, A)$$

Definition 1.4 (Capture-free substitution).

A substitution σ is *capture-free* for a formula A if for every free variable x in A , none of the variables in $\sigma(x)$ is bound in A .

Note: if $t \in \mathcal{T}$ is a closed term, then $\{y \setminus t\}$ is capture-free for any A .

Outline

- ▶ Preliminaries and Reminders
- ▶ **Soundness Proof**
- ▶ Completeness: Preliminaries
- ▶ Proof of Completeness
- ▶ Examples of Counter-model Construction

How to show the Soundness Theorem?

As for propositional logic, we show the following lemmas:

1. All LK-rules preserve falsifiability upwards.
2. An LK-derivation with a falsifiable root sequent has at least one falsifiable leaf sequent
3. All axioms are valid

Finally, we use these lemmas to show the soundness theorem.

Preservation of Falsifiability

Definition 2.1.

An LK-rule θ *preserves falsifiability* (upwards) if whenever the conclusion w of an instance $\frac{w_1 \cdots w_n}{w}$ of θ is falsifiable, then also at least one of the premises w_i is falsifiable

NEW: the falsifying interpretation for the conclusion does not need to be the same as for the conclusion.

Lemma 2.1.

All LK-rules preserve falsifiability.

- ▶ We have shown that the rules for propositional connectives (\wedge , \vee , \rightarrow , \neg) have this property.
- ▶ It remains to show that also the \forall and \exists rules preserve falsifiability.

Proof: \forall -left preserves falsifiability

$$\frac{\Gamma, \forall x A, A[x \setminus t] \implies \Delta}{\Gamma, \forall x A \implies \Delta} \forall\text{-left} \quad t \text{ is a closed term}$$

- ▶ Assume that $\mathcal{I} = (D, \iota)$ falsifies the conclusion $\Gamma, \forall x A \implies \Delta$.
- ▶ \mathcal{I} makes all formulae in $\Gamma \cup \{\forall x A\}$ true and all formulae in Δ false.
- ▶ It suffices to show that $\mathcal{I} \models A[x \setminus t]$. Then, the premiss is falsified by \mathcal{I} .
- ▶ Since $\mathcal{I} \models \forall x A$, we know that $v_{\mathcal{I}}(\alpha\{x \leftarrow d\}, A) = T$ for all $d \in D$ and any α . (Using the semantics of \forall)
- ▶ In particular, $v_{\mathcal{I}}(\alpha\{x \leftarrow v_{\mathcal{I}}(\alpha, t)\}, A) = T$
- ▶ By the substitution lemma: $v_{\mathcal{I}}(\alpha, A[x \setminus t]) = T$
- ▶ And therefore: $\mathcal{I} \models A[x \setminus t]$.

Proof: \exists -left preserves falsifiability

$$\frac{\Gamma, A[x \setminus a] \implies \Delta}{\Gamma, \exists x A \implies \Delta} \exists\text{-left} \quad a \text{ is a constant that does not occur in the conclusion}$$

- ▶ Assume that $\mathcal{I} = (D, \iota)$ falsifies the conclusion $\Gamma, \exists x A \implies \Delta$.
- ▶ \mathcal{I} makes all formulae in $\Gamma \cup \{\exists x A\}$ true and all formulae in Δ false.
- ▶ We have to find an interpretation that falsifies the premiss.
- ▶ But we can **not** simply assume that $\mathcal{I} \models A[x \setminus a]$.
- ▶ Since $\mathcal{I} \models \exists x A$ we know that $v_{\mathcal{I}}(\alpha\{x \leftarrow d\}, A) = T$ for some $d \in D$.
- ▶ Based on \mathcal{I} and d , we define a **new** model \mathcal{I}' as follows:
 - ▶ $\mathcal{I}' = (D, \iota')$ is identical to \mathcal{I} except for the interpretation of the constant a .
 - ▶ a is interpreted as d , ie. $a^{\iota'} = d$.
- ▶ Then \mathcal{I}' falsifies the premiss:
 - ▶ Since a does not occur in the conclusion, \mathcal{I}' and \mathcal{I} interpret the fml. in Γ and Δ equally. \mathcal{I}' makes all fml. in Γ true and all fml. in Δ false.
 - ▶ $v_{\mathcal{I}'}(\alpha\{x \leftarrow d\}, A) = T$, and $d = v_{\mathcal{I}'}(\alpha, a)$, so $\mathcal{I}' \models A[x \setminus a]$.

An Example

- ▶ Assume that $\mathcal{I} = (D, \iota)$ is an interpretation with domain $D = \{1, 2\}$ and $p^\iota = \{2\}$.
- ▶ Assume that a og b are constants and $a^\iota = b^\iota = 1$.
- ▶ Then $\mathcal{I} \not\models p(a)$ og $\mathcal{I} \not\models p(b)$.

$$\frac{p(b) \implies p(a)}{\exists x p(x) \implies p(a)} \exists\text{-left}$$

- ▶ \mathcal{I} falsifies the conclusion:

$$\begin{aligned} \mathcal{I} &\models \exists x p(x), \text{ since } v_{\mathcal{I}}(\alpha\{x \leftarrow 2\}, p(x)) = T \\ \mathcal{I} &\not\models p(a). \end{aligned}$$

- ▶ But \mathcal{I} does not falsify the premiss because $\mathcal{I} \not\models p(b)$.
- ▶ We define a new interpretation $\mathcal{I}' = (D, \iota')$ such that $b^{\iota'} = 2$.
- ▶ Then \mathcal{I}' falsifies the premiss.

Proof: \exists -right and \forall -right preserve satisfiability

- ▶ The proof for \forall -right is dual to that for \exists -left
- ▶ The proof for \exists -right is dual to that for \forall -left

How to show the Soundness Theorem?

As for propositional logic, we show the following lemmas:

1. All LK-rules preserve falsifiability upwards.
2. An LK-derivation with a falsifiable root sequent has at least one falsifiable leaf sequent
3. All axioms are valid

Finally, we use these lemmas to show the soundness theorem.

Existence of a falsifiable leaf sequent

Lemma 2.2.

If the root sequent \mathcal{I} of an LK-derivation is falsifiable, then at least one of the leaf sequents is satisfiable.

- ▶ As for propositional logic, the proof is by structural induction on the LK-derivation.
- ▶ The base case (one sequent $\Gamma \Longrightarrow \Delta$) is trivial since $\Gamma \Longrightarrow \Delta$ is both root and leaf sequent.
- ▶ Two induction steps, for one-premise and two-premise rules
- ▶ Both use the lemma that falsifiability is preserved upwards.

Difference from propositional logic: not necessarily the same interpretation!

How to show the Soundness Theorem?

As for propositional logic, we show the following lemmas:

1. All LK-rules preserve falsifiability upwards.
2. An LK-derivation with a falsifiable root sequent has at least one falsifiable leaf sequent
3. All axioms are valid

Finally, we use these lemmas to show the soundness theorem.

All axioms are valid

Lemma 2.3.

All axioms are valid

- ▶ The proof is the same as for propositional logic
- ▶ An axiom has the form

$$\Gamma, p(t_1, \dots, t_n) \implies p(t_1, \dots, t_n), \Delta$$

- ▶ Any interpretation that satisfies the antecedent satisfies $p(t_1, \dots, t_n)$.
- ▶ Therefore, the same formula $p(t_1, \dots, t_n)$ is satisfied in the succedent.

Proof of the Soundness Theorem for LK

Proof of soundness.

- ▶ Assume that \mathcal{P} is an LK-proof for the sequent $\Gamma \Longrightarrow \Delta$.
 - ▶ \mathcal{P} is an LK-derivation where every leaf is an axiom.
- ▶ For the sake of contradiction, assume that $\Gamma \Longrightarrow \Delta$ is **not** valid.
- ▶ Then there is a countermodel \mathcal{I} that falsifies $\Gamma \Longrightarrow \Delta$.
- ▶ We know from the previous Lemma that there is an \mathcal{I}' that falsifies at least one leaf sequent of \mathcal{P} .
- ▶ Then \mathcal{P} has a leaf sequent that is not an axiom, since axioms are not falsifiable.
- ▶ So \mathcal{P} cannot be an LK-proof.



Outline

- ▶ Preliminaries and Reminders
- ▶ Soundness Proof
- ▶ **Completeness: Preliminaries**
- ▶ Proof of Completeness
- ▶ Examples of Counter-model Construction

Herbrand Universe

Definition 3.1 (Herbrand universe).

Let T be a set of terms. Then $\mathcal{H}(T)$, the *Herbrand universe of T* , is the smallest set such that

- ▶ $\mathcal{H}(T)$ contains all constant symbols from T . If there are no constants in T , we include some constant symbol o from \mathcal{A} (called a dummy constant) in $\mathcal{H}(T)$.
- ▶ If f is a function symbol in T , with arity n and t_1, \dots, t_n are terms in $\mathcal{H}(T)$, then $f(t_1, \dots, t_n) \in \mathcal{H}(T)$.

The Herbrand universe of a set of formulae is the Herbrand universe of the set of terms occurring in the formulae. The Herbrand universe of a branch of a derivation is the Herbrand universe of the set of formulae occurring on that branch.

- ▶ Intuitively, the Herbrand universe of T is the set of all *closed* terms that can be constructed from the constant and function symbols in T .

Herbrand Universe: Examples

Example.

Let $T = \{f(x)\}$. Then the Herbrand universe of T is the set

$$\{o, f(o), f(f(o)), f(f(f(o))), \dots\}$$

Example.

Let $T = \{a, f(x)\}$. Then the Herbrand universe of T is the set

$$\{a, f(a), f(f(a)), f(f(f(a))), \dots\}$$

Example.

Let $F = \{\forall x p(f(g(x)))\}$. Then the Herbrand universe of F is the set

$$\{o, f(o), g(o), f(g(o)), g(f(o)), f(f(o)), g(g(o)), \dots\}$$

Fairness

- ▶ To guarantee that a proof is found
 - ▶ all formulae have to be used in a rule eventually, and
 - ▶ all \forall -left and \exists -right rules are applied with *all terms* eventually.
- ▶ If we try to guarantee this,
 1. Either all branches can be closed, giving a proof,
 2. or there is an open branch that we can generate a counterexample from.
- ▶ This only makes sense if we include infinite derivations, i.e. derivations with infinitely long branches.
- ▶ We construct a *limit* by either continuing until no more rules can be applied, or continuing to apply rules indefinitely. We call the result of this process a **limit derivation**.
- ▶ When we talk about limit derivations, we include infinite trees.
- ▶ We won't define these formally.
- ▶ If all branches in a derivation can be closed, then the derivation is finite. I.e. proofs are finite.

Fairness

Definition 3.2 (Fair derivations).

A limit derivation is *fair* if each open branch has the following properties:

1. There are no sequents $\Gamma, A \implies A, \Delta$ on the branch that could be closed using the axiom.
2. If a \wedge , \vee , \rightarrow , or \neg formula occurs, then the corresponding LK rule is applied to the formula on that branch.
3. If a \exists formula occurs in an antecedent, or a \forall formula in a succedent, then the \exists -left, resp. \forall -right rules are applied to the formula on that branch.
4. If a \forall formula occurs in an antecedent, or a \exists formula in a succedent, then the \forall -left, resp. \exists -right rules are applied to the formula on that branch *for every term t* in the Herbrand universe of that branch.

Königs Lemma

Lemma 3.1 (Königs lemma).

If T is an infinite tree, but finitely branching (all nodes have finitely many descendants), then T has an infinitely long branch.

Proof.

We inductively define an infinitely long branch. Let u_0 be the root node of the tree T . Since T is infinite and u_0 has finitely many descendants, one of u_0 's descendants must be infinite. (Otherwise T would be finite.) Let u_1 be the root of such a sub-tree. If the branch u_0, u_1, \dots, u_n is defined, we find the next node u_{n+1} by the same kind of reasoning. This process defines an infinitely long branch. □

Corollary 3.1.

If T is a finitely branching tree, where all branches are finitely long, then T is finite.

Outline

- ▶ Preliminaries and Reminders
- ▶ Soundness Proof
- ▶ Completeness: Preliminaries
- ▶ **Proof of Completeness**
- ▶ Examples of Counter-model Construction

Proof of Completeness

Assume $\Gamma \implies \Delta$ is not provable.

- ▶ Construct a fair (limit) derivation \mathcal{D} from $\Gamma \implies \Delta$. Possibly infinite.
- ▶ Then there is (at least) one branch \mathcal{B} that does not end in an axiom.
- ▶ We construct an interpretation that falsifies $\Gamma \implies \Delta$. Let

\mathcal{B}^\top be the set of formulae that occur in an antecedent on \mathcal{B} , and
 \mathcal{B}^\perp be the set of formulae that occur in a succedent on \mathcal{B} , and
 At be the set of *atomic* formulae in \mathcal{B}^\top .

Proof of Completeness (Construction of counter-model)

- ▶ We construct a counter-model $\mathcal{I} = (D, \iota)$ for $\Gamma \implies \Delta$.
- ▶ Let the domain D be the Herbrand universe of the branch. (I.e. the set of all closed terms that can be generated from the terms on the branch).
- ▶ Let $a^\iota = a$ for all constant symbols $a \in \mathcal{A}$.
- ▶ If $f \in \mathcal{F}$ is a function symbol with arity n , let $f^\iota(t_1, \dots, t_n) = f(t_1, \dots, t_n)$.
 - ▶ Then $v_{\mathcal{I}}(t) = t$ for all closed terms t .
 - ▶ All terms are interpreted as themselves
- ▶ If p is a predicate symbol with arity n , let $\langle t_1, \dots, t_n \rangle \in p^\iota$ if and only if $p(t_1, \dots, t_n) \in \Delta$.
- ▶ Such an interpretation is often called a **Herbrand model** or a **term model**.

Proof of Completeness (Properties of \mathcal{I})

- ▶ We show by structural induction on first-order formulae that the interpretation \mathcal{I} makes *all* formulae in \mathcal{B}^\top true and all formulae in \mathcal{B}^\perp false.
- ▶ We show for all first-order formulae A that:
 - If $A \in \mathcal{B}^\top$, then $\mathcal{I} \models A$, i.e. $v_{\mathcal{I}}(A) = T$
 - If $A \in \mathcal{B}^\perp$, then $\mathcal{I} \not\models A$, i.e. $v_{\mathcal{I}}(A) = F$

Base case 1: A is an atomic formula $p(t_1, \dots, t_n)$ in \mathcal{B}^\top .

- ▶ Then $p(t_1, \dots, t_n) \in \text{At}$ og $\langle t_1, \dots, t_n \rangle \in p^v$ by construction.
- ▶ Therefore $\mathcal{I} \models p(t_1, \dots, t_n)$.

Base case 2: A is an atomic formula $p(t_1, \dots, t_n)$ in \mathcal{B}^\perp .

- ▶ Since \mathcal{B} does not end in an axiom, and the derivation is fair, $p(t_1, \dots, t_n) \notin \text{At}$ and $\langle t_1, \dots, t_n \rangle \notin p^v$.
- ▶ Therefore $\mathcal{I} \not\models p(t_1, \dots, t_n)$.

Proof of Completeness (Propositional connectives)

Induction step: From the assumption (induction hypothesis) that our statement holds for all smaller formulae, we have to show that it holds for $\neg A$, $(A \wedge B)$, $(A \vee B)$, $(A \rightarrow B)$, $\forall x A$, and $\exists x A$.

Most of this was done in the proof for propositional logic

E.g. assume that $A \wedge B \in \mathcal{B}^\top$.

- ▶ By fairness of the derivation, the \wedge -left rule has been applied to $A \wedge B$ on the branch \mathcal{B} .
- ▶ Then $A \in \mathcal{B}^\top$ and $B \in \mathcal{B}^\top$.
- ▶ By the induction hypothesis, $\mathcal{I} \models A$ and $\mathcal{I} \models B$.
- ▶ By model semantics, $\mathcal{I} \models A \wedge B$.

We only need to cover quantified formulae

Proof of Completeness (\exists in Antecedent)

Assume that $\exists x A \in \mathcal{B}^\top$.

- ▶ By fairness of the derivation, \exists -left was applied to $\exists x A$ on the branch.
- ▶ Then there is a constant a such that $A[x \setminus a] \in \mathcal{B}^\top$.
- ▶ By the ind. hyp., $\mathcal{I} \models A[x \setminus a]$.
- ▶ I.e. $v_{\mathcal{I}}(\alpha, A[x \setminus a]) = T$ for any assignment α , since $A[x \setminus a]$ is closed
- ▶ By the substitution lemma: $v_{\mathcal{I}}(\alpha \{x \leftarrow a^t\}, A) = T$.
- ▶ By model semantics: $v_{\mathcal{I}}(\alpha, \exists x A) = T$
- ▶ I.e. $\mathcal{I} \models \exists x A$.

Proof of Completeness (\exists in Succedent)

Assume that $\exists x A \in \mathcal{B}^\perp$.

- ▶ We have to show that $\mathcal{I} \not\models \exists x A$. Assume that this does not hold.
- ▶ I.e. $\mathcal{I} \models \exists x A$
- ▶ Remember that the domain D of $\mathcal{I} = (D, \iota)$ consists of terms
- ▶ Then $v_{\mathcal{I}}(\alpha\{x \leftarrow t\}, A) = T$ for some term $t \in D$.
- ▶ By fairness of the derivation, the \exists -right rule was applied on $\exists x A$ with the term t .
- ▶ It follows that:
 - ▶ $A[x \setminus t] \in \mathcal{B}^\perp$
 - ▶ $v_{\mathcal{I}}(A[x \setminus t]) = F$ (induction hypothesis)
 - ▶ $v_{\mathcal{I}}(\alpha\{x \leftarrow v_{\mathcal{I}}(t)\}, A) = F$ for any α (substitution lemma)
 - ▶ $v_{\mathcal{I}}(\alpha\{x \leftarrow t\}, A) = F$ (since $v_{\mathcal{I}}(t) = t$)
- ▶ Contradiction!

Proof of Completeness (\forall in Succedent)

Assume that $\forall x A \in \mathcal{B}^\perp$.

- ▶ By fairness of the derivation, \forall -right was applied to $\exists x A$ on the branch.
- ▶ Then there is a constant a such that $A[x \setminus a] \in \mathcal{B}^\perp$.
- ▶ By the ind. hyp., $\mathcal{I} \not\models A[x \setminus a]$.
- ▶ I.e. $v_{\mathcal{I}}(\alpha, A[x \setminus a]) = F$ for any assignment α , since $A[x \setminus a]$ is closed
- ▶ By the substitution lemma: $v_{\mathcal{I}}(\alpha \{x \leftarrow a^l\}, A) = F$.
- ▶ By model semantics: $v_{\mathcal{I}}(\alpha, \forall x A) = F$
- ▶ I.e. $\mathcal{I} \not\models \forall x A$.

Proof of Completeness (\forall in Antecedent)

Assume that $\forall x A \in \mathcal{B}^\top$.

- ▶ We have to show that $\mathcal{I} \models \forall x A$. Assume that this does not hold.
- ▶ I.e. $\mathcal{I} \not\models \forall x A$
- ▶ Remember that the domain D of $\mathcal{I} = (D, \iota)$ consists of terms
- ▶ Then $v_{\mathcal{I}}(\alpha\{x \leftarrow t\}, A) = F$ for some term $t \in D$.
- ▶ By fairness of the derivation, the \forall -left rule was applied on $\forall x A$ with the term t .
- ▶ It follows that:
 - ▶ $A[x \setminus t] \in \mathcal{B}^\top$
 - ▶ $v_{\mathcal{I}}(A[x \setminus t]) = T$ (induction hypothesis)
 - ▶ $v_{\mathcal{I}}(\alpha\{x \leftarrow v_{\mathcal{I}}(t)\}, A) = T$ for any α (substitution lemma)
 - ▶ $v_{\mathcal{I}}(\alpha\{x \leftarrow t\}, A) = T$ (since $v_{\mathcal{I}}(t) = t$)
- ▶ Contradiction!

Some comments

- ▶ We can see the construction of a limit derivation as approximating a counter-model for $\Gamma \implies \Delta$.
- ▶ The more often we apply the \forall -left and \exists -right rules, the 'closer' we get to a possible counter-model
- ▶ But constructing a counter-model in this way may require using all rules infinitely often.
- ▶ So this is **not an algorithm** for finding counter-models!
- ▶ It will find a proof if one exists, but may not terminate otherwise.
- ▶ There may be finite counter-models even when this method does not terminate. Finding finite counter-models is a topic of active research.
- ▶ The idea of the completeness proof is important: we construct an interpretation from something purely syntactic.

Outline

- ▶ Preliminaries and Reminders
- ▶ Soundness Proof
- ▶ Completeness: Preliminaries
- ▶ Proof of Completeness
- ▶ **Examples of Counter-model Construction**

Counter-model Construction, Ex. 1

$$\begin{array}{c}
 \mathcal{B} \\
 \hline
 qa, A, pa \Rightarrow qb, pb \quad \quad \quad \overline{qb, qa, A, pa \Rightarrow qb} \\
 \hline
 qa, A, pb \rightarrow qb, pa \Rightarrow qb \\
 \hline
 qa, A, pa \Rightarrow qb \\
 \hline
 qa, A, pa \Rightarrow \forall x qx \\
 \hline
 \overline{A, pa \Rightarrow \forall x qx, pa} \\
 \hline
 \underbrace{\frac{A, pa \rightarrow qa, pa \Rightarrow \forall x qx}{\forall x (px \rightarrow qx), pa \Rightarrow \forall x qx}}_A
 \end{array}$$

- ▶ Abbreviate px for $p(x)$, qb for $q(b)$, etc.
- ▶ The Herbrand universe of branch \mathcal{B} , and domain of \mathcal{I} , is $\{a, b\}$.
- ▶ Since $pa \in \mathcal{B}^\top$, define $a \in p^\iota$, so $\mathcal{I} \models pa$.
- ▶ Since $qa \in \mathcal{B}^\top$, define $a \in q^\iota$, so $\mathcal{I} \models qa$ and thus $\mathcal{I} \models pa \rightarrow qa$.
- ▶ Since $qb \in \mathcal{B}^\perp$, define $b \notin q^\iota$, so $\mathcal{I} \not\models qb$ and thus $\mathcal{I} \not\models \forall x qx$.
- ▶ Since $pb \in \mathcal{B}^\perp$, define $b \notin p^\iota$, so $\mathcal{I} \not\models pb$ and thus $\mathcal{I} \models pb \rightarrow qb$.
- ▶ Therefore also $\mathcal{I} \models \forall x (px \rightarrow qx)$.
- ▶ \mathcal{I} makes all of \mathcal{B}^\top true and all of \mathcal{B}^\perp false.

Summary and Outlook

- ▶ We can show things for ∞ many interpretations using finite proofs!
- ▶ OMG! Amazing!
- ▶ Uncloseable branches give counter-models
- ▶ Might be infinite: if there is no proof, we might search for ever
- ▶ First-order validity is undecidable

- ▶ Can this be automated?
- ▶ Sure! But...
- ▶ Instantiating quantifiers with *every* possible term is wasteful
- ▶ More goal-oriented ways of doing this?
- ▶ Coming up...