

Mandag 7. september

1.

Rotasjon om (1,1)
med vinkel π

1 Flytt $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ til origo

2 Rotasjon med π

3 Flytt tilbake

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \left(\begin{pmatrix} x \\ y \end{pmatrix} - \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right) + \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$= \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 2 \\ 2 \end{pmatrix}$$

$$\text{dvs } \underline{\underline{T_{\begin{pmatrix} 2 \\ 2 \end{pmatrix}} S_{\pi}}}$$

Utleddning av trigonometriske likheter

$$S_{\varphi} S_{\psi} = S_{\varphi+\psi} \quad \begin{pmatrix} \cos \varphi & -\sin \varphi \\ \sin \varphi & \cos \varphi \end{pmatrix} \begin{pmatrix} \cos \psi & -\sin \psi \\ \sin \psi & \cos \psi \end{pmatrix} = \begin{pmatrix} \cos(\varphi+\psi) & -\sin(\varphi+\psi) \\ \sin(\varphi+\psi) & \cos(\varphi+\psi) \end{pmatrix}$$

$$\cos \varphi \cos \psi - \sin \varphi \sin \psi = \cos(\varphi+\psi)$$

$$\sin \varphi \cos \psi + \cos \varphi \sin \psi = \sin(\varphi+\psi)$$

Refleksjon over linja $x - 2y = 2$

$$\text{Normalvektor } \begin{pmatrix} 1 \\ -2 \end{pmatrix} = \vec{v}$$

$$\text{Flytter linja til gjennom origo} \quad \begin{pmatrix} x \\ y \end{pmatrix} - \begin{pmatrix} 0 \\ -1 \end{pmatrix} = \begin{pmatrix} x' \\ y' \end{pmatrix}$$

$$x' - 2y' = x - 2(y+1) = x - 2y - 2 = 2 - 2 = 0$$

$$Q_{\vec{v}} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - 2 \frac{1}{5} \begin{pmatrix} 1 & -2 \\ -2 & 4 \end{pmatrix} = \frac{1}{5} \begin{pmatrix} 3 & 4 \\ 4 & -3 \end{pmatrix}$$

$$\frac{1}{5} \begin{pmatrix} 3 & 4 \\ 4 & -3 \end{pmatrix} \left(\begin{pmatrix} x \\ y \end{pmatrix} - \begin{pmatrix} 0 \\ -1 \end{pmatrix} \right) + \begin{pmatrix} 0 \\ -1 \end{pmatrix} = \frac{1}{5} \begin{pmatrix} 3 & 4 \\ 4 & -3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} \frac{4}{5} \\ -\frac{3}{5} \end{pmatrix} + \begin{pmatrix} 0 \\ -1 \end{pmatrix}$$

$$= \frac{1}{5} \begin{pmatrix} 3 & 4 \\ 4 & -3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \frac{1}{5} \begin{pmatrix} 4 \\ -8 \end{pmatrix}$$

F.eks. $x=4, y=1$:

$$\frac{1}{5} \begin{pmatrix} 3 & 4 \\ 4 & -3 \end{pmatrix} \begin{pmatrix} 4 \\ 1 \end{pmatrix} + \frac{1}{5} \begin{pmatrix} 4 \\ -8 \end{pmatrix} = \frac{1}{5} \left(\begin{pmatrix} 16 \\ 13 \end{pmatrix} + \begin{pmatrix} 4 \\ -8 \end{pmatrix} \right) = \begin{pmatrix} 4 \\ 1 \end{pmatrix}$$

$$L_G = \{t_{\bar{a}} \in G\}$$

Diskret undergruppe av $\text{Isom}_2 \cong G$ ²

$$= \{t_{m\bar{a}+n\bar{b}}; m, n \in \mathbb{Z}\}$$

$$\forall t_{\bar{a}} \in G, \bar{a} \neq 0, \|\bar{a}\| \geq \varepsilon$$

$$\forall S_{\theta, p} \in G, \theta \neq 0, |\theta| \geq \varepsilon$$

Gitter av rang 2

Mengden av gitterpunkt $M_G = \{m\bar{a} + n\bar{b} \mid m, n \in \mathbb{Z}\}$

$\forall$ kein ante $\bar{a}, \bar{b} \geq 0$ - (spuss)

Dirichlet område:

$$D_G = \{p \in \mathbb{R}^2 \mid d(p, 0) \leq d(p, q) \forall q \in M_G, q \neq 0\}$$



Punktgruppe av $t_{\bar{a}} T: T$

Setning: Punktgrupper for D_G pøsses selv.

$$g \in t_{\bar{a}} T \quad g t_{\bar{a}} g^{-1} = t_{\bar{b}} T t_{\bar{a}} T^{-1} t_{-\bar{b}}$$

$$= t_{\bar{b}} t_{T(\bar{a})} T T^{-1} t_{-\bar{b}}$$

$$T t_{\bar{a}} = t_{T(\bar{a})} T$$

$$= t_{T(\bar{a})}$$

ford.

$$T(x + \bar{a}) = T(x) + T(\bar{a})$$


$$d(T(\bar{x}), 0) = d(T(\bar{x}), T(0)) = d(\bar{x}, \bar{0})$$

$$T(M_G) = M_G \quad d(T(\bar{x}), \bar{y}) = d(T(\bar{x}), T(\bar{z})) = d(\bar{x}, \bar{z}) \quad \forall \bar{x}, \bar{y}, \bar{z} \in M_G \quad (\neq \bar{0})$$

$$\Rightarrow \{p \in \mathbb{R}^2 \mid d(T(p), 0) \leq d(p, q)\} = \{p \in \mathbb{R}^2 \mid d(p, 0) \leq d(p, q)\}$$

Setning Punktgrupper til en diskret undergruppe
av Isom_2 med $\text{rg } L_G = 2$ er en av C_n, D_n
men $n = 1, 2, 3, 4, 6$

Skisse av bevis



Rotasjon med vinkel $\theta = \frac{2\pi}{n}$

$$d = \sqrt{(\cos \theta - 1)^2 + (\sin \theta)^2}$$

$$= \sqrt{\cos^2 \theta - 2\cos \theta + 1 + \sin^2 \theta}$$

$$= \sqrt{2 - 2\cos \theta} \geq 1 \quad \theta \neq 0$$

altså $2 - 2\cos \theta \geq 1$

$$\Rightarrow \cos \theta \leq \frac{1}{2}$$

$$\theta = \frac{2\pi}{n} \Rightarrow \frac{2\pi}{n} \geq \frac{2\pi}{6}$$

$$\Rightarrow n \leq 6$$

altså $n = 1, 2, 3, 4, 5, 6$

Men $n=5$ gir



$$\| (\cos \frac{4\pi}{5}, \sin \frac{4\pi}{5}) + (1, 0) \|^2 = (\cos \frac{4\pi}{5} + 1)^2 + \sin^2 \frac{4\pi}{5}$$

$$= 2 + 2\cos \frac{4\pi}{5} = 2 - 2\cos \frac{2\pi}{5}$$

$$= 2 + 2(2\cos^2 \frac{2\pi}{5} - 1) = 4\cos^2 \frac{2\pi}{5}$$

$$= 4 \cdot \left(\frac{\sqrt{5}-1}{4}\right)^2 = \frac{1}{4}(5 - 2\sqrt{5} + 1) = \frac{3-\sqrt{5}}{2} < 1$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta$$

$$= \cos^2 \theta - 1 + \cos^2 \theta$$

$$= 2\cos^2 \theta - 1$$