

Nilpotence and Stable Homotopy Theory II

Author(s): Michael J. Hopkins and Jeffrey H. Smith

Source: *Annals of Mathematics*, Second Series, Vol. 148, No. 1 (Jul., 1998), pp. 1-49

Published by: Mathematics Department, Princeton University

Stable URL: <https://www.jstor.org/stable/120991>

Accessed: 02-10-2018 09:56 UTC

REFERENCES

Linked references are available on JSTOR for this article:

https://www.jstor.org/stable/120991?seq=1&cid=pdf-reference#references_tab_contents

You may need to log in to JSTOR to access the linked references.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



JSTOR

Mathematics Department, Princeton University is collaborating with JSTOR to digitize, preserve and extend access to *Annals of Mathematics*

Nilpotence and stable homotopy theory II

By MICHAEL J. HOPKINS* and JEFFREY H. SMITH*

Contents

Introduction

1. Morava K-theories
 - 1.1. Construction
 - 1.2. Spectra related to BP
 - 1.3. Fields in the category of spectra
 - 1.4. Morava K-theories and duality
 - 1.5. Proof of Corollary 5, assuming Theorem 3
 2. Proofs of Theorems 3 and 7
 - 2.1. Proof of Theorem 3
 - 2.2. Proof of Theorem 7
 3. v_n self-maps
 4. Construction of v_n self-maps
 - 4.1. Preliminaries
 - 4.2. Vanishing lines
 - 4.3. Morava K-theories and the Adams spectral sequence
 - 4.4. Examples of self maps
 - 4.5. Proof that Theorem 7 implies Theorem 2
 5. Endomorphisms, up to nilpotents
 - 5.1. N -endomorphisms
 - 5.2. Classification of N -endomorphisms
 - 5.3. Some technical tools
 - 5.4. N -endomorphisms and thick subcategories
 - 5.5. A spectrum with few nonnilpotent self maps
 - 5.6. Proof of Theorem 5.3
- Appendix A. Proof of Theorem 4.13
- References

*Partially supported by the NSF.

Introduction

This paper is a continuation of [11]. Since so much time has elapsed since its publication, a recasting of the context is in order.

In [19] Ravenel described a series of conjectures getting at the structure of stable homotopy theory in the large. The theory was organized around a family of “higher periodicities” generalizing Bott periodicity, and it depended on being able to determine the nilpotent and non-nilpotent maps in the category of spectra. There are three senses in which a map of spectra can be nilpotent:

Definition 1. i) A map of spectra $f: F \rightarrow X$ is *smash nilpotent* if for $n \gg 0$ the map $f^{(n)}: F^{(n)} \rightarrow X^{(n)}$ is null.

ii) A self-map $f: \Sigma^k F \rightarrow F$ is *nilpotent* if for $n \gg 0$ the map $f^n: \Sigma^{kn} F \rightarrow F$ is null.

iii) A map $f: S^m \rightarrow R$ from the sphere spectrum to a ring spectrum is *nilpotent* if it is nilpotent when regarded as an element of the ring $\pi_* R$.

The following is the main result of [11].

THEOREM 2 ([11, Th. 1]). *In each of the above situations, the map f is nilpotent if the spectrum F is finite and the map $1_{MU} \wedge f$ is null. In cases ii) and iii) it suffices to assume that $MU_* f = 0$.*

In case the range of f is p -local, the condition can be replaced with the condition $1_{BP} \wedge f$ (resp. $BP_* f = 0$).

The purpose of this paper is to refine this criterion and to produce some interesting nonnilpotent maps. Many of the results of this paper were conjectured by Ravenel in [19].

Let $K(n)$ be the n^{th} Morava K-theory at the prime p (see §1).

THEOREM 3. i) *Let R be a (p -local) ring spectrum. An element $\alpha \in \pi_* R$ is nilpotent if and only if for all $0 \leq n \leq \infty$, $K(n)_*(\alpha)$ is nilpotent.*

ii) *A self-map $f: \Sigma^k F \rightarrow F$, of the p -localization of a finite spectrum, is nilpotent if and only if $K(n)_* f$ is nilpotent for all $0 \leq n < \infty$.*

iii) *A map $f: F \rightarrow X$ from a finite spectrum to a p -local spectrum is smash nilpotent if and only if $K(n)_* f = 0$ for all $0 \leq n \leq \infty$.*

Of course, the hypothesis “ p -local” can be dropped if the condition on the Morava K theory is checked at all primes.

Theorem 3 can be used to determine which cohomology theories detect the non-nilpotent maps in the category of spectra.

Definition 4. A ring spectrum E is said to *detect nilpotence* if, equivalently,
 i) for any ring spectrum R , the kernel of the Hurewicz homomorphism $E_*: \pi_*R \rightarrow E_*R$ consists of nilpotent elements;
 ii) a map $f: F \rightarrow X$ from a finite spectrum F to any spectrum X is smash nilpotent if $1_E \wedge f: E \wedge F \rightarrow E \wedge X$ is null homotopic.

To see that the parts of this definition are equivalent, replace MU by E in the remark on page 209 of [11] and in the proof on page 211 of [11] that Theorem 2 implies part ii) of Theorem 1.

COROLLARY 5. *A ring spectrum E detects nilpotence if and only if*

$$K(n)_*E \neq 0$$

for all $0 \leq n \leq \infty$ and for all primes p .

Now let \mathcal{C}_0 be the homotopy category of p -local finite spectra, let $\mathcal{C}_n \subset \mathcal{C}_0$ be the full subcategory of $K(n-1)$ -acyclics, and let \mathcal{C}_∞ be the full subcategory consisting of contractible spectra. The \mathcal{C}_n fit into a sequence

$$\mathcal{C}_\infty \subset \cdots \subset \mathcal{C}_{n+1} \subset \mathcal{C}_n \subset \cdots \subset \mathcal{C}_0.$$

This is a nontrivial fact. That there are inclusions $\mathcal{C}_{n+1} \subset \mathcal{C}_n$ is essentially the Invariant Prime Ideal Theorem (see [19]). That the inclusions are proper is a result of Steve Mitchell [17].

Definition 6. A full subcategory \mathcal{C} of the category of spectra is said to be *thick* if it is closed under weak equivalences, cofiber sequences and retracts; i.e.,

- i) An object weakly equivalent to an object of \mathcal{C} is in \mathcal{C} .
- ii) If $X \rightarrow Y \rightarrow Z$ is a cofiber sequence, and two of $\{X, Y, Z\}$ are in \mathcal{C} then so is the third.
- iii) A retract of an object of \mathcal{C} is in \mathcal{C} .

THEOREM 7. *If $\mathcal{C} \subseteq \mathcal{C}_0$ is a thick subcategory, then $\mathcal{C} = \mathcal{C}_n$ for some n .*

Theorem 7 is in fact equivalent to Theorem 2 (the proof is sketched at the end of Section 4). It is often used in the following manner.

Call a property P of p -local finite spectra *generic* if the full subcategory of \mathcal{C}_0 consisting of the objects satisfying P is thick. To show that $X \in \mathcal{C}_n$ has a generic property P it suffices (by Theorem 7) to show that *any* object of $\mathcal{C}_n \setminus \mathcal{C}_{n+1}$ has P . The proofs of the next few results use this technique.

Theorem 3 limits the non-nilpotent maps in \mathcal{C}_0 —they must be detected by some Morava K -theory. The simplest type is a v_n self-map.

Definition 8. Let X be a p -local finite spectrum, and $n \geq 0$. A self-map $v: \Sigma^k X \rightarrow X$ is said to be a v_n -self-map if

$$K(m)_*v \text{ is } \begin{cases} \text{multiplication by a} & \text{if } m = n = 0; \\ \text{rational number} & \\ \text{an isomorphism} & \text{if } m = n \neq 0; \\ \text{nilpotent} & \text{if } m \neq n. \end{cases}$$

It turns out that the property of admitting a v_n self-map is generic.

THEOREM 9. *A p -local finite spectrum X admits a v_n self-map if and only if $X \in C_n$. If X admits a v_n self-map, then for $N \gg 0$, X admits a v_n self-map*

$$v: \Sigma^{p^N 2(p^n-1)} X \rightarrow X$$

satisfying

$$(*) \quad K(m)_*v = \begin{cases} v_n^{p^N} & \text{if } m = n; \\ 0 & \text{otherwise.} \end{cases}$$

The v_n self-maps turn out to be distinguished by another property.

Definition 10. A ring homomorphism

$$f: A \rightarrow B$$

is an F -isomorphism if

- i) the kernel of f consists of nilpotent elements, and
- ii) given $b \in B$, b^{p^n} is in the image of f for some n .

Two rings A and B are F -isomorphic ($A \approx_F B$) if there is an F -isomorphism between them.

THEOREM 11. *Let $X \in C_n \setminus C_{n+1}$. The $K(n)$ -Hurewicz homomorphism gives rise to an F -isomorphism*

$$(12) \quad \text{center } [X, X]_* \approx_F \begin{cases} \mathbb{Z}_{(p)} & (n = 0) \\ \mathbf{F}_p[v_n] & (n \neq 0). \end{cases}$$

Put another way, this result shows that the v_n maps essentially constitute the center of the endomorphism rings of finite spectra.

The description of spectra as cell complexes encourages the intuition that the endomorphism rings of finite spectra approximate matrix algebras over the ring $\pi_* S^0$. This would suggest that the centers of these rings are generated by the maps obtained by smashing the identity map with a map between spheres—an impossibility by Theorem 11. A more accurate description might be that the ‘Morita’ equivalence classes of these rings are determined by the integer n

of Theorem 11. This integer invariant can also be thought of as determining the ‘birational’ equivalence classes of finite spectra. For more on this analogy see [13].

There is a less metaphorical interpretation of the integer which occurs in Theorems 9 and 11. It turns out to correspond to the Bousfield class ([19, 1.19],[9]) of the spectrum X .

Definition 13. Two spectra X and Y are *Bousfield equivalent* if they have the same acyclic spectra:

$$\text{For all } Z, \quad X \wedge Z \sim * \Leftrightarrow Y \wedge Z \sim *.$$

The *Bousfield class* of X (denoted $\langle X \rangle$) is the collection of spectra Y which are Bousfield equivalent to X

There is a natural partial ordering on Bousfield classes of spectra. One writes $\langle X \rangle \geq \langle Y \rangle$ if for all Z ,

$$X \wedge Z \sim * \Rightarrow Y \wedge Z \sim *.$$

Thus the class sphere spectrum is maximal with respect to this ordering, and the class of contractible spectra is minimal. There are also two binary operations (\vee and \wedge). They are defined by

$$\begin{aligned} \langle X \rangle \vee \langle Y \rangle &= \langle X \vee Y \rangle, \\ \langle X \rangle \wedge \langle Y \rangle &= \langle X \wedge Y \rangle. \end{aligned}$$

The operation \wedge distributes over \vee .

Let \mathbf{N} be the set of nonnegative integers and \mathbf{P} the set of primes. For a finite spectrum X , let $\text{Cl}(X) \subseteq \mathbf{N} \times \mathbf{P}$ denote the set of pairs (n, p) for which $K(n)_*X \neq 0$ at p .

THEOREM 14. *If X and Y are finite spectra, then $\langle X \rangle \leq \langle Y \rangle$ if and only if $\text{Cl}(X) \subseteq \text{Cl}(Y)$.*

Theorem 14 affirms Ravenel’s class invariance conjecture ([19]).

Proof of Theorem 14. It is immediate from the definition that if $\langle X \rangle \leq \langle Y \rangle$ then $\text{Cl}(X) \subseteq \text{Cl}(Y)$. Suppose then that $\text{Cl}(X) \subseteq \text{Cl}(Y)$: Since $\langle X \rangle \leq \langle Y \rangle$ if and only if $\langle X_{(p)} \rangle \leq \langle Y_{(p)} \rangle$ for all prime p , we may localize everything at a prime p . Having done that, suppose that $Y \in \mathcal{C}_n \setminus \mathcal{C}_{n+1}$. By assumption, X is also in \mathcal{C}_n . For a fixed Y , the property (of X)

$$\langle X \rangle \leq \langle Y \rangle$$

is a generic property. It follows that the class

$$\{ X \mid \langle X \rangle \leq \langle Y \rangle \}$$

is equal to \mathcal{C}_m for some m . Since $\langle Y \rangle = \langle Y \rangle$, $m \leq n$ (in fact $m = n$), and so $X \in \mathcal{C}_m$. This completes the proof. \square

The organization of this paper is as follows. In Section 1 the Morava K-theories and related spectra are introduced, and their basic properties are established. Section 2 contains the proofs of Theorems 3 and 7. The section ends with a simple argument reducing Corollary 5 to Theorem 3. In Section 3, v_n self-maps are introduced and their basic properties are established. The main result of the section is that the property of admitting a v_n self-map is generic. This reduces the existence of v_n self-maps to the task of constructing a single example of each type. These examples are constructed in Section 4. The construction makes use of the Adams spectral sequence. Section 5 is concerned with the classification of the endomorphisms, up to nilpotent elements, of full subcategories which are stable under suspension, of the homotopy category of finite spectra. The proof of Theorem 11 is given after Corollary 5.4. The appendix to the paper contains some results on the cohomology of Hopf algebras which are needed in Section 4.

Acknowledgments. Most of the results of this paper date from 1985, and many people have helped shape their course. Special thanks are due to Emmanuel Dror-Farjoun whose prodding eventually led to the formulation of Theorem 7, and to Clarence Wilkerson for helpful conversations concerning the proof of Theorem 4.13. Even deeper debts are owed to Doug Ravenel for formulating such a beautiful body of conjectures, and to Mark Mahowald for placing in the hands of the authors the tools for proving results like these. Finally, the first author would like to dedicate his contributions to this paper to Ruth, Randi and Rose.

Notation and conventions. We will work for the most part in the homotopy category of spectra as defined in [2], [14], or [10], and not in some model category of spectra. This is in part because the main results of this paper concern the homotopy category. It also allows us to avoid choosing a particular model.

This expository decision comes at a mild cost. For instance, in order to form the mapping cone of a map $f: X \rightarrow Y$ in the homotopy category, one must represent it by a map

$$\tilde{f}: \tilde{X} \rightarrow \tilde{Y}$$

in some model of the category of spectra, and then define $Y \cup_f CX$ to be the image in the homotopy category of the mapping cone

$$\tilde{Y} \cup_f C\tilde{X} = \tilde{Y} \amalg \tilde{X} \times I / \sim$$

of \tilde{f} . The isomorphism (weak equivalence) class of $Y \cup_f CX$ depends only on the original map f , but its dependence on f is not functorial. Nevertheless,

this object will be referred to as the *mapping cone of f*, or as the *cofiber of f*, and will be denoted C_f .

Having chosen a lift of \tilde{f} of f , one has a natural identification of the mapping cone of $\tilde{Y} \rightarrow \tilde{Y} \cup \tilde{X}$ with $\tilde{X} \wedge S^1$, and this latter map is used to define the connecting homomorphism in the long exact sequence of a cofibration. Because if this, we will take the assertion that a sequence

$$X \xrightarrow{f} Y \rightarrow Z$$

is a *cofiber sequence* to mean that it comes equipped with a map $Z \rightarrow X \wedge S^1$, and that there is a map $\tilde{f}: \tilde{X} \rightarrow \tilde{Y}$ of cofibrant objects in some model of the category of spectra, with the property that the image of

$$\tilde{X} \rightarrow \tilde{Y} \rightarrow \tilde{Y} \cup \tilde{X} \rightarrow \tilde{X} \wedge S^1$$

becomes isomorphic to

$$X \rightarrow Y \rightarrow Z \rightarrow X \wedge S^1$$

in the homotopy category. With this convention, enough structure has been specified to give all of the usual long exact sequences associated to a cofiber sequence.

A *map of cofiber sequences*

$$(X_1 \rightarrow Y_1 \rightarrow Z_1) \rightarrow (X_2 \rightarrow Y_2 \rightarrow Z_2)$$

consists of maps

$$X_1 \xrightarrow{f} X_2, \quad Y_1 \rightarrow Y_2, \quad \text{and} \quad Z_1 \rightarrow Z_2$$

for which the following diagram commutes:

$$\begin{array}{ccccccc} X_1 & \longrightarrow & Y_1 & \longrightarrow & Z_1 & \longrightarrow & X_1 \wedge S^1 \\ f \downarrow & & \downarrow & & \downarrow & & \downarrow f \wedge S^1 \\ X_2 & \longrightarrow & Y_2 & \longrightarrow & Z_2 & \longrightarrow & X_2 \wedge S^1. \end{array}$$

A map of cofiber sequences gives rise to maps between all of the usual long exact sequences.

As described above, any map $X \rightarrow Y$ can be extended to a cofiber sequence. Given cofiber sequences

$$X_1 \xrightarrow{i_1} Y_1 \rightarrow Z_1 \quad \text{and} \quad X_2 \xrightarrow{i_2} Y_2 \rightarrow Z_2,$$

and commutative square

$$\begin{array}{ccc} X_1 & \xrightarrow{i_1} & Y_1 \\ \downarrow & & \downarrow \\ X_1 & \xrightarrow{i_2} & Y_1, \end{array}$$

there always exists a map $Z_1 \rightarrow Z_2$ extending the square to a map of cofiber sequences. This map, however, is not unique.

All of this basically amounts to saying that the homotopy category of spectra is a *triangulated category* with the cofiber sequences as triangles.

A square

$$\begin{array}{ccc} W & \longrightarrow & X \\ \downarrow & & \downarrow \\ Y & \longrightarrow & Z \end{array}$$

is *homotopy cocartesian* if any of the following equivalent conditions holds:

(1) There is a lift of the square to some model of the category of spectra of the form

$$\begin{array}{ccc} \tilde{W} & \xrightarrow{i} & \tilde{X} \\ j \downarrow & & \downarrow \\ \tilde{Y} & \longrightarrow & \tilde{Z}, \end{array}$$

with W , X , Y , and Z cofibrant, the maps i and j cofibrations, and the map

$$X \cup_W Y \rightarrow Z$$

a weak equivalence.

(2) There is an extension of the square to a map of cofiber sequences

$$\begin{array}{ccccc} W & \longrightarrow & X & \longrightarrow & A \\ \downarrow & & \downarrow & & \downarrow \\ Y & \longrightarrow & Z & \longrightarrow & B \end{array}$$

in which the map $A \rightarrow B$ is a weak equivalence.

(3) In any extension of the square to a map of cofiber sequences

$$\begin{array}{ccccc} W & \longrightarrow & X & \longrightarrow & A \\ \downarrow & & \downarrow & & \downarrow \\ Y & \longrightarrow & Z & \longrightarrow & B, \end{array}$$

the map $A \rightarrow B$ is a weak equivalence.

A homotopy cocartesian square gives rise to a family of Meyer-Vietoris sequences.

The reader may have noticed that we have chosen to form the cofiber of a map $f: X \rightarrow Y$ of spectra with the “cone coordinate” on the right. With this convention the canonical weak equivalence of the cofiber of

$$Z \wedge X \rightarrow Z \wedge Y$$

with $Z \wedge (Y \cup_f CX)$ involves only the associativity of the smash product, and not the process of moving the cone coordinate past Z . Also with this convention, the cofiber of $Y \rightarrow Y \cup_f CX$ is naturally identified with $X \wedge S^1$ (and not $S^1 \wedge X$). This avoids the troublesome sign that can crop up when one tries to relate the connecting homomorphism in a cofiber sequence with the connecting homomorphism in some suspension of the cofiber sequence.

A sequence

$$X_0 \rightarrow X_1 \rightarrow \dots$$

in the homotopy category of spectra can always be lifted to a sequence of cofibrations between cofibrant objects

(15)
$$\tilde{X}_0 \rightarrow \tilde{X}_1 \rightarrow \dots$$

in some model for the category of spectra. The image of

$$\tilde{X}_\infty = \varinjlim \tilde{X}_i$$

in the homotopy category is independent of the choice of lift, and depends functorially on the sequence (15). We will write $\varinjlim X_i$ for the image of \tilde{X}_∞ in the homotopy category of spectra, and refer to it as the *homotopy colimit* of the sequence (15). The homotopy colimit is rarely the colimit because of the \varprojlim^1 -term in the Milnor sequence.

The assumption that a spectrum is finite is made several times. In contexts when the category in mind is the category of p -local spectra, this term is used to refer to a spectrum which is weakly equivalent to the p -localization of a finite spectrum. The only property of finite spectra used is that the set of homotopy classes of maps from a finite spectrum to a sequential homotopy colimit is the colimit of the maps

$$[X, \varinjlim Y_\alpha] = \varinjlim [X, Y_\alpha].$$

In general, an object of a category with this property is said to be *small*. It can be shown that the small objects of the category of p -local spectra are precisely the objects which are weakly equivalent to the p -localizations of a finite spectrum.

A spectrum X is *connective* if $\pi_k X = 0$ for $k \ll 0$. It is *connected* if $\pi_k X = 0$ for $k < 0$. Thus “connected” and “ (-1) -connected” are synonymous. Similarly, a graded abelian group is *connective* if the homogeneous part of degree k is zero for $k \ll 0$. A graded abelian group is *connected* if the homogeneous component of degree k is zero for $k < 0$.

The Eilenberg-MacLane spectrum with coefficients in an abelian group A will be denoted HA . To be consistent with this, the homology of a spectrum X with coefficients in A will be denoted HA_*X .

Finally, the suspension of a map will always be labeled with the same symbol as the map.

1. Morava K-theories

1.1. *Construction.* The study of a ring is often simplified by passage to its quotients and localizations. The same is true of ring *spectra*, though constructing quotients and localizations can be difficult. In good cases the following constructions can be made:

Quotients. Suppose that E is a ring spectrum and that $\pi_*E = R$ is commutative. Given $x \in R$, define the spectrum $E/(x)$ by the cofiber sequence

$$\Sigma^{|x|}E \xrightarrow{x} E \rightarrow E/(x).$$

If x is not a zero divisor then $\pi_*E/(x)$ is isomorphic to the ring $R/(x)$. In good cases $E/(x)$ will still be a ring spectrum, and the map

$$E \rightarrow E/(x)$$

will be a map of ring spectra. Given a regular sequence

$$\{x_1, \dots, x_n, \dots\} \subset R,$$

one can hope to iterate the above construction and form a ring spectrum

$$E/(x_1, \dots, x_n, \dots)$$

with

$$\pi_*E/(x_1, \dots, x_n, \dots) \approx R/(x_1, \dots, x_n, \dots),$$

and such that the natural map

$$E \rightarrow E/(x_1, \dots, x_n, \dots)$$

is a map of ring spectra.

Localizations. Let E and R be as above, and suppose that $S \subset R$ is a multiplicatively closed subset. Since $S^{-1}R$ is a flat R -module, the functor

$$S^{-1}E_*(_) \stackrel{\text{def}}{=} S^{-1}R \otimes_R E_*(_)$$

is a homology theory. In good cases it is represented by a ring spectrum, and the localization map by a map of ring spectra

$$E \rightarrow S^{-1}E.$$

1.2. *Spectra related to BP.* When the ring spectrum in question is BP , the above constructions can always be made, using the Baas-Sullivan theory of bordism with singularities. See [5], [18], [22] for the details.

Recall that $BP_* \approx \mathbb{Z}_{(p)}[v_1, \dots, v_n \dots]$ with $|v_n| = 2p^n - 2$. To fix notation, take the set $\{v_n\}$ to be the Hazewinkel generators [12]. For $0 < n < \infty$ the ring spectra $K(n)$ and $P(n)$ are defined by the isomorphisms

$$\begin{aligned} K(n)_* &\approx \mathbb{F}_p[v_n, v_n^{-1}], \\ P(n)_* &\approx \mathbb{F}_p[v_n, v_{n+1} \dots], \\ P(0)_* &\approx BP_*, \end{aligned}$$

with the understanding that they are constructed from BP using a combination of the above methods. It is also useful to set

$$\begin{aligned} K(0) &= H\mathbb{Q}, \\ K(\infty) &= H\mathbb{F}_p. \end{aligned}$$

There are maps $P(n) \rightarrow P(n + 1)$, and the colimit

$$\lim_n P(n)$$

is the Eilenberg-MacLane spectrum $H\mathbb{F}_p$.

PROPOSITION 1.1 ([19, Th. 2.1(c)]). *The Bousfield classes of $K(n)$ and $P(n)$ are related by*

$$\langle P(n) \rangle = \langle K(n) \rangle \vee \langle P(n + 1) \rangle.$$

Consequently,

$$\langle BP \rangle = \langle K(0) \rangle \vee \dots \vee \langle K(n) \rangle \vee \langle P(n + 1) \rangle.$$

Proof. The proposition follows from the next two results of Ravenel [19]. □

PROPOSITION 1.2 ([19, Lemma 1.34]). *Let $v: \Sigma^k X \rightarrow X$ be a self-map of a spectrum X . Let X/vX and $v^{-1}X$ denote the cofiber of v and the homotopy colimit of the sequence*

$$\dots \Sigma^{-k|v|} X \xrightarrow{v} \Sigma^{-(k+1)|v|} X \rightarrow \dots$$

respectively. Then there is an equality of Bousfield classes

$$\langle X \rangle = \langle X/vX \rangle \vee \langle v^{-1}X \rangle. \quad \square$$

PROPOSITION 1.3 ([19, Th. 2.1(a)]). *There is an equality of Bousfield classes*

$$\langle v_n^{-1}P(n) \rangle = \langle K(n) \rangle. \quad \square$$

1.3. *Fields in the category of spectra.* The coefficient ring $K(n)_*$ is a graded field in the sense that all of its graded modules are free. This begets a host of special properties of the Morava K-theories.

PROPOSITION 1.4. *For any spectrum X , $K(n) \wedge X$ has the homotopy type of a wedge of suspensions of $K(n)$.*

Proof. Choose a basis $\{e_i\}_{i \in I}$ of the free $K(n)_*$ -module $K(n)_*X$, and represent it as a map

$$\bigvee_{i \in I} S^{|e_i|} \rightarrow K(n) \wedge X.$$

The composition

$$K(n) \wedge \bigvee_{i \in I} S^{|e_i|} \rightarrow K(n) \wedge K(n) \wedge X \rightarrow K(n) \wedge X$$

is then a weak equivalence. \square

PROPOSITION 1.5. *For any two spectra X and Y , the natural map*

$$(1.6) \quad K(n)_*X \otimes_{K(n)_*} K(n)_*Y \rightarrow K(n)_*X \wedge Y$$

is an equivalence.

Proof. Consider the map (1.6) as a transformation of functors of Y . The left side satisfies the Eilenberg-Steenrod axioms since $K(n)_*Y$ is a flat (in fact free) $K(n)_*$ -module. The right side satisfies the Eilenberg-Steenrod axioms by definition. The transformation is an isomorphism when Y is the sphere, hence for all Y . \square

Propositions 1.4 and 1.5 portray the Morava K-theories as being a lot like fields. One formulation of Theorem 3 is that they are the *prime fields* of the category of spectra.

A (skew) field is a ring, all of whose modules are free.

Definition 1.7. A non-contractible ring spectrum E is a *field* if E_*X is a free E_* -module for all spectra X .

This property also admits a geometric expression.

LEMMA 1.8. *If E is a field, then $E \wedge X$ has the homotopy type of a wedge of suspensions of E .*

Proof. This is very similar to the proof of 1.4. \square

PROPOSITION 1.9. *Let E be a field. Then E has the homotopy type of a wedge of suspensions of $K(n)$ for some n .*

Proof. Since $1 \in \pi_*E$ is non-nilpotent, Theorem 3 implies that for some prime p and for some $n \leq \infty$,

$$K(n)_*E \neq 0.$$

Since $K(n)$ and E are both fields, it follows from Lemma 1.8 that $K(n) \wedge E$ is both a wedge of suspensions of $K(n)$ and a wedge of suspensions of E . In particular, E is a retract of a wedge of suspensions of $K(n)$. The result therefore follows from the next proposition. \square

PROPOSITION 1.10. *Let M have the homotopy type of a wedge of suspensions of $K(n)$ (fixed n). If E is a retract of M , then E itself has the homotopy type of a wedge of suspensions of $K(n)$.*

LEMMA 1.11. *The homotopy homomorphism induced by the Hurewicz map*

$$\iota \wedge 1_M: M \approx S^0 \wedge M \rightarrow K(n) \wedge M$$

is a homomorphism of $K(n)_$ -modules.*

Proof. The map in question is a wedge of suspensions of the map

$$\eta_R: K(n) \approx S^0 \wedge K(n) \rightarrow K(n) \wedge K(n),$$

so it suffices to prove the claim when M is $K(n)$. In this case the result is a consequence of the formula [20, A.2.2.5 or 6.1.13],

$$\eta_R(v_n) = v_n. \quad \square$$

LEMMA 1.12. *Let $f: M \rightarrow N$ be a map of wedges of suspensions of $K(n)$. The homotopy homomorphism*

$$\pi_*f: \pi_*M \rightarrow \pi_*N$$

is a map of $K(n)_$ -modules.*

Proof. Consider the following commutative diagram:

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \downarrow & & \downarrow \\ K(n) \wedge M & \xrightarrow{1 \wedge f} & K(n) \wedge N. \end{array}$$

The right vertical arrow is the inclusion of a wedge summand since N admits the structure of a $K(n)$ -module spectrum. It therefore suffices to prove that the composition induces a map of $K(n)_*$ -modules. The left vertical arrow does by Lemma 1.11 and the bottom horizontal arrow is a map of $K(n)$ -module spectra. \square

Proof of Proposition 1.10. Since M has the homotopy type of a wedge of suspensions of $K(n)$'s, it can be given the structure of a $K(n)$ -module spectrum. Let $i: E \rightarrow M$, and $p: M \rightarrow E$ be the inclusion and retraction mappings respectively. By Lemma 1.12, the composite $i \circ p$ induces a homomorphism of $K(n)_*$ -modules

$$\pi_*M \rightarrow \pi_*M.$$

Choose a basis $\{e_i\}$ of the image of this map, and represent it by

$$\bigvee S^{|e_i|} \rightarrow M.$$

The map

$$N = K(n) \wedge \left(\bigvee S^{|e_i|}\right) \rightarrow K(n) \wedge M \rightarrow M$$

then gives rise to an isomorphism

$$\pi_*N \approx \text{image of } \pi_*(i \circ p),$$

since it sends the obvious basis of π_*N to the basis $\{e_i\}$. The composite

$$N \rightarrow M \xrightarrow{p} E$$

is the desired homotopy equivalence. □

1.4. *Morava K-theories and duality.* We will often use the device of replacing a self map of a finite spectrum

$$f: \Sigma^n X \rightarrow X$$

with its Spanier-Whitehead dual

$$Df: S^n \rightarrow X \wedge DX,$$

a map from the n -sphere to the *ring spectrum* $X \wedge DX$. If V is the finite-dimensional $K(n)_*$ -vector space $K(n)_*X$, then the ring $K(n)_*(X \wedge DX)$ is naturally isomorphic to the ring

$$V \otimes V^* \approx \text{End}(V).$$

The effect in Morava K-theory of the duality map

$$X \wedge DX \xrightarrow{\text{flip}} DX \wedge X \xrightarrow{\text{duality}} S^0$$

is to send an endomorphism to its trace (in the graded sense). Let $\{e_i\} \subset V$ be a basis of V , and $\{e_i^*\} \subset V^*$ the corresponding dual basis. The effect of the other duality map

$$S^0 \rightarrow X \wedge DX$$

is to send $1 \in K(n)_*$ to $\sum e_i \otimes e_i^* \in V \otimes V^*$. In particular,

LEMMA 1.13. *The duality map $S^0 \rightarrow X \wedge DX$ induces a nonzero homomorphism in $K(n)$ -homology if and only if $K(n)_*X \neq 0$.*

1.5. *Proof of Corollary 5, assuming Theorem 3.* Let R be a ring spectrum, and $\alpha \in \pi_*R$. By Proposition 1.4, if $K(n)_*E \neq 0$, then

$$E_*\alpha = 0 \Rightarrow K(n)_*\alpha = 0.$$

It follows that if for all n , $K(n)_*E \neq 0$, then for all n , $K(n)_*\alpha = 0$, so that α is nilpotent by Theorem 3. On the other hand, if $K(n)_*E = 0$ for some n , then E does not detect the non-nilpotent map

$$\iota: S^0 \rightarrow K(n),$$

so E does not detect nilpotence. This completes the proof.

2. Proofs of Theorems 3 and 7

2.1. *Proof of Theorem 3.* Some of the conditions in Theorem 3 require the case $n = \infty$, and some of them do not. When the target spectrum is finite, the case $n = \infty$ is superfluous.

LEMMA 2.1. *Let X and Y be finite spectra. For $m \gg 0$,*

- i) $K(m)_*X \approx H\mathbb{F}_{p_*}X \otimes K(m)_*$;
- ii) $K(m)_*Y \approx H\mathbb{F}_{p_*}Y \otimes K(m)_*$;
- iii) $K(m)_*f = H\mathbb{F}_{p_*}f \otimes 1_{K(m)_*}$ for every $f: X \rightarrow Y$.

Proof. This follows from the Atiyah-Hirzebruch spectral sequence, when we use the fact that $|v_m| \rightarrow \infty$ as $m \rightarrow \infty$. □

COROLLARY 2.2. *If f is either a self-map of a finite spectrum or an element in the homotopy of a finite ring spectrum, the following are equivalent:*

- i) $K(m)_*f$ is nilpotent for $m \gg 0$;
- ii) $H\mathbb{F}_{p_*}f$ is nilpotent.

If $|f| \neq 0$ then both of these conditions hold.

Proof. If $|f| \neq 0$ then, from dimensional considerations,

$$H\mathbb{F}_{p_*}f^i = 0 \quad \text{for } i \gg 0.$$

It then follows from 2.1 that $K(m)_*f^i = 0$ for $i, m \gg 0$. When $|f| = 0$, part (3) of 2.1 applies to every power of f . The result follows easily from this. □

Let $f: S^0 \rightarrow X$ be a map of spectra. Consider the homotopy direct limit T of the sequence

$$(2.3) \quad S^0 \rightarrow X \rightarrow X \wedge X \rightarrow X \wedge X \wedge X \rightarrow \dots,$$

in which the map $X^{(n)} \rightarrow X^{(n+1)}$ is given by

$$f \wedge 1_{X^{(n)}}: X^{(n)} \approx S^0 \wedge X^{(n)} \rightarrow X^{(n+1)}.$$

The n -fold composition

$$S^0 \rightarrow \dots \rightarrow X^{(n)}$$

is the iterated smash product

$$f^{(n)} = f \wedge \dots \wedge f.$$

The map

$$f^{(\infty)}: S^0 \rightarrow T$$

can be thought of as the *infinite smash product* of f .

LEMMA 2.4. *Let E be a ring spectrum with unit $\iota: S^0 \rightarrow E$. The following are equivalent:*

- i) $E \wedge T$ is contractible;
- ii) $\iota \wedge f^{(\infty)}: S^0 \rightarrow E \wedge T$ is null;
- iii) $\iota \wedge f^{(n)}: S^0 \rightarrow E \wedge X^{(n)}$ is null for $n \gg 0$;
- iv) $1_E \wedge f^{(n)}: E \approx E \wedge S^0 \rightarrow E \wedge X^{(n)}$ is null for $n \gg 0$.

Proof. i) \Rightarrow ii) and iv) \Rightarrow i) are immediate. Since

$$\varinjlim E \wedge X^{(n)} \approx E \wedge T,$$

and since homotopy groups commute with direct limits, a null homotopy of $S^0 \rightarrow E \wedge T$ must occur at some $S^0 \rightarrow E \wedge X^{(n)}$ for $n \gg 0$. This gives ii) \Rightarrow iii). The implication iii) \Rightarrow iv) is the only one requiring E to be a ring spectrum. If $S^0 \rightarrow E \wedge X^{(n)}$ is null then so is the first map in the following factorization of $1_E \wedge f^{(n)}$:

$$E \wedge S^0 \rightarrow E \wedge E \wedge X^{(n)} \rightarrow E \wedge X^{(n)}.$$

This completes the proof. □

Proof of Theorem 3. Part i) follows from part iii), since the iterated multiplication factors through iterated smashing. Part ii) follows from part i) since multiplication in the rings

$$\pi_* X \wedge DX \quad \text{and} \quad K(n)_* X \wedge DX$$

corresponds, under Spanier-Whitehead duality, to composition in

$$[X, X]_* \quad \text{and} \quad \text{End}_{K(n)_*}(K(n)_* X).$$

Replacing

$$f: F \rightarrow X$$

with

$$Df: S^0 \rightarrow DF \wedge X$$

in part iii) if necessary, we may assume that $F = S^0$. The result reduces to Theorem 2 once it is shown that

$$1_{BP} \wedge f^{(m)}$$

is null for $m \gg 0$. From Lemma 2.4 (with the obvious notation) this is equivalent to showing that $BP \wedge T$ is contractible. In view of the Bousfield equivalence

$$\langle BP \rangle = \langle K(0) \rangle \vee \cdots \vee \langle K(n) \rangle \vee \langle P(n+1) \rangle,$$

it is enough to show that $P(n) \wedge T$ is contractible for $n \gg 0$. Again from 2.4 this is equivalent to showing that

$$S^0 \rightarrow P(n) \wedge T$$

is null for $n \gg 0$. Now let n grow to infinity. Since

$$\varinjlim P(n) \approx H\mathbb{F}_p,$$

the map

$$S^0 \rightarrow \varinjlim P(n) \wedge T$$

is null by assumption. Since homotopy commutes with direct limits, the null homotopy arises at some

$$S^0 \rightarrow P(n) \wedge T.$$

This completes the proof of Theorem 3. □

2.2. *Proof of Theorem 7.* The proof of Theorem 7 requires a slight modification of the third assertion of Theorem 3, and a useful cofiber sequence (2.7).

COROLLARY 2.5. *Let F and Z be finite spectra, E a ring spectrum, and X an arbitrary spectrum.*

i) *If a map $f: F \rightarrow X \wedge E$ satisfies*

$$K(n)_*(f) = 0 \quad \text{for all } 0 \leq n \leq \infty,$$

then for $m \gg 0$, the composite

$$F^{(m)} \xrightarrow{f^{(m)}} (X \wedge E)^{(m)} \approx X^{(m)} \wedge E^{(m)} \xrightarrow{1 \wedge \mu} X^{(m)} \wedge E$$

is null.

ii) A map

$$f: F \rightarrow X$$

has the property that

$$f^{(m)} \wedge 1_Z: F^{(m)} \wedge Z \rightarrow X^{(m)} \wedge Z$$

is null for $m \gg 0$ if and only if

$$K(n)_*(f \wedge 1_Z) = 0$$

for all $0 \leq n \leq \infty$.

Proof. In part i), the map $f^{(m)}$ is already null for $m \gg 0$ by part iii) of Theorem 3. The *only if* part of ii) is clear. Letting E be the ring spectrum $Z \wedge DZ$ and replacing

$$f \wedge 1_Z: F \wedge Z \rightarrow X \wedge Z$$

with its Spanier-Whitehead dual

$$F \rightarrow X \wedge Z \wedge DZ$$

reduce the *if* part to i). □

LEMMA 2.6. Let $X \xrightarrow{f} Y \xrightarrow{g} Z$ be a sequence of maps. The map $C_f \rightarrow C_{g \circ f}$ induced by g gives rise to a cofiber sequence

$$C_f \rightarrow C_{g \circ f} \rightarrow C_g.$$

Proof. Consider the following diagram in which the rows and columns are cofiber sequences:

$$\begin{array}{ccccc}
 X & \xrightarrow{f} & Y & \longrightarrow & C_f \\
 \parallel & & g \downarrow & & \downarrow \\
 X & \xrightarrow{g \circ f} & Z & \longrightarrow & C_{g \circ f} \\
 & & \downarrow & & \downarrow \\
 & & C_g & \longrightarrow & ?
 \end{array}$$

The upper right square is a pushout. It follows that the bottom arrow is a homotopy equivalence. This completes the proof. □

COROLLARY 2.7. Let $f: X \rightarrow Y$ and $g: Z \rightarrow W$ be two maps. There is a cofiber sequence

$$X \wedge C_g \rightarrow C_{f \wedge g} \rightarrow C_f \wedge Y.$$

Proof. Apply the lemma to the factorization

$$f \wedge g = f \wedge 1_Y \circ 1_X \wedge g. \quad \square$$

Proof of Theorem 7. It suffices to establish the following:

$$(2.8) \quad \text{if } X \in \mathcal{C} \text{ and } X \in \mathcal{C}_n \text{ then } \mathcal{C}_n \subseteq \mathcal{C},$$

for then $\mathcal{C} = \mathcal{C}_m$, where

$$m = \min \{ n \mid \mathcal{C}_n \subseteq \mathcal{C} \}.$$

Since everything has been localized at p , set

$$\text{Cl}(X) = \{ n \in \mathbf{N} \mid K(n)_*(X) \neq 0 \}.$$

With this notation, (2.8) becomes:

$$(2.9) \quad \text{if } X \in \mathcal{C} \text{ and } \text{Cl}(Y) \subset \text{Cl}(X), \text{ then } Y \in \mathcal{C}.$$

Suppose, then, that $X \in \mathcal{C}$. Then so is $Z \wedge X$ for any $Z \in \mathcal{C}_0$. Let $f: F \rightarrow S^0$ be the fiber of the duality map $S^0 \rightarrow X \wedge DX$. Then $Y \wedge C_f \in \mathcal{C}$. Setting $g = f^{(m-1)}$ in Corollary 2.7 and smashing with the identity map of Y gives a cofiber sequence

$$Y \wedge F \wedge C_{f^{(m-1)}} \rightarrow Y \wedge C_{f^{(m)}} \rightarrow Y \wedge C_f \wedge F^{(m-1)}.$$

It follows that $Y \wedge C_{f^{(m)}} \in \mathcal{C}$ for all m .

By 1.13, $K(n)_*f \neq 0$ if and only if $n \notin \text{Cl}(X)$, so that

$$K(n)_*(1_Y \wedge f) = 0 \quad \text{for all } n,$$

since $\text{Cl}(Y) \subseteq \text{Cl}(X)$. Part ii) of Corollary 2.5 then gives that $1_Y \wedge f^{(m)}$ is null for $m \gg 0$. This means that

$$Y \wedge C_{f^{(m)}} \approx Y \vee (\Sigma Y \wedge F^{(m)}) \quad \text{for } m \gg 0,$$

so that $Y \in \mathcal{C}$. This completes the proof of Theorem 7. □

3. v_n Self-maps

The purpose of the next two sections is to establish Theorem 9. The “only if” part, that $X \notin \mathcal{C}_n$ implies that X does not admit a v_n self-map is easy: if for some $j < n$, $K(j)_*X \neq 0$, and if v is a v_n self-map, then the cofiber Y of v is a finite spectrum satisfying

$$K(n)_*Y = 0, \quad K(j)_*Y \neq 0,$$

contradicting the fact that $\mathcal{C}_n \subset \mathcal{C}_j$. The proof of the “if” part falls into two steps. In this section it is shown that the property of admitting a v_n self-map

is generic. It then remains to construct for each n , a spectrum X_n with a v_n self-map. This is done in Section 4.

For any spectrum X , the element $p \in [X, X]_*$ is a v_0 self-map satisfying condition (*) of Theorem 9. We therefore need only consider v_n self-maps when $n \geq 1$. Because of this, unless otherwise mentioned, *in this and the next section, we will work entirely in the category \mathcal{C}_1 .*

As mentioned in subsection 1.4, a self-map

$$\Sigma^k F \rightarrow F$$

of a finite spectrum corresponds, under Spanier-Whitehead duality, to a map from the k -sphere to the ring spectrum

$$R = F \wedge DF.$$

Definition 3.1. Let R be a finite ring spectrum, $n > 0$. An element

$$\alpha \in \pi_* R$$

is a v_n -element if

$$K(m)_* \alpha \text{ is } \begin{cases} \text{a unit} & \text{if } m = n \\ \text{nilpotent} & \text{otherwise.} \end{cases}$$

LEMMA 3.2. Let R be a finite ring spectrum, and $\alpha \in \pi_* R$ a v_n -element. There exist integers i and j such that

$$K(m)_* \alpha^i = \begin{cases} 0 & \text{if } m \neq n \\ v_n^j & \text{if } m = n. \end{cases}$$

Proof. It follows from Corollary 2.2 that $H\mathbb{F}_{p_*} \alpha$ is nilpotent. Raising α to a power, if necessary, we may suppose that $H\mathbb{F}_{p_*} \alpha = 0$. It then follows from Corollary 2.2 that $K(m)_* \alpha = 0$ for all but finitely many m . Raising α to a further power, if necessary, it can be arranged that $K(m)_* \alpha = 0$ for $m \neq n$.

The assertion $K(n)_* \alpha^i = v_n^j$ is equivalent to the assertion that $\alpha^i = 1 \in K(n)_* R / (v_n - 1)$. The ring $K(n)_* R / (v_n - 1)$ has a finite group of units, so i can be taken to be the order of this group. □

COROLLARY 3.3. If $f: \Sigma^k F \rightarrow F$ is a v_n self-map, then there exist integers i, j with the property that

$$K(m)_* f^i = \begin{cases} 0 & \text{if } m \neq n \\ \text{multiplication by } v_n^j & \text{if } m = n. \end{cases}$$

LEMMA 3.4. Suppose that x and y are commuting elements of a $\mathbb{Z}_{(p)}$ -algebra. If $x - y$ is both torsion and nilpotent, then for $N \gg 0$,

$$x^{p^N} = y^{p^N}.$$

Proof. Since we are working over a $\mathbb{Z}_{(p)}$ -algebra it follows that

$$p^k(x - y) = 0$$

for some k . The result now follows by expansion of

$$x^{p^N} = (y + (x - y))^{p^N}$$

using the binomial theorem. □

LEMMA 3.5. *Let R be a finite ring spectrum, and $\alpha \in \pi_k R$ a v_n -element. For some $i > 0$, α^i is in the center of $\pi_* R$.*

Proof. Raising α to a power, if necessary, we may assume that $K(m)_* \alpha$ is in the center of $K(m)_* R$ for all m . Let $l(\alpha)$ and $r(\alpha)$ be the elements of $\text{End}(\pi_* R)$ given by left and right multiplication by α . Since $R \in \mathcal{C}_1$ the difference $l(\alpha) - r(\alpha)$ has finite order. Since

$$K(m)_* (l(\alpha) - r(\alpha)) = 0 \quad \text{for all } m,$$

$l(\alpha) - r(\alpha)$ is nilpotent by Theorem 3. The result now follows from 3.4 (with $x = l(\alpha)$, $y = r(\alpha)$). □

LEMMA 3.6. *Let $\alpha, \beta \in \pi_* R$ be v_n -elements. There exist integers i and j with $\alpha^i = \beta^j$.*

Proof. Raising α and β to powers if necessary, we may assume that for all m , $K(m)_* (\alpha - \beta) = 0$. The result follows, as above, from 3.4. □

COROLLARY 3.7. *If f and g are two v_n self-maps of F , then f^i is homotopic to g^j for some i and j .*

COROLLARY 3.8. *Suppose X and Y have v_n self-maps v_X and v_Y . There are integers i and j so that for every Z and every*

$$f: Z \wedge X \rightarrow Y$$

the following diagram commutes:

$$\begin{array}{ccc} Z \wedge X & \xrightarrow{f} & Y \\ 1 \wedge v_X^i \downarrow & & \downarrow v_Y^j \\ Z \wedge X & \xrightarrow{f} & Y. \end{array}$$

Proof. The spectrum $DX \wedge Y$ has two v_n self-maps: $Dv_X \wedge 1_Y$ and $1_{DX} \wedge v_Y$. By Corollary 3.7 there are integers i and j for which $Dv_X^i \wedge 1_Y$ is homotopic to $1_{DX} \wedge v_Y^j$. The result now follows from Spanier-Whitehead duality. □

COROLLARY 3.9. *The full subcategory of \mathcal{C}_1 consisting of spectra admitting a v_n self-map is thick.*

Proof. Call the subcategory in question \mathcal{C} . Note that $X \in \mathcal{C}$ if and only if $\Sigma X \in \mathcal{C}$. To check that \mathcal{C} is closed under cofiber sequences it therefore suffices to show that if

$$(3.10) \quad X \rightarrow Y \rightarrow Z$$

is a cofiber sequence with X and Y in \mathcal{C} , then Z is in \mathcal{C} . Using Corollary 3.8 choose the v_n self-maps v_X and v_Y of X and Y so that

$$\begin{array}{ccccc} \Sigma^k X & \longrightarrow & \Sigma^k Y & \longrightarrow & \Sigma^k Z \\ v_X \downarrow & & v_Y \downarrow & & \\ X & \longrightarrow & Y & \longrightarrow & Z \end{array}$$

commutes. It is easy to check that any map $v_Z: \Sigma^k Z \rightarrow Z$ making the above diagram a map of cofiber sequences is a v_n self-map.

Now suppose that Y is a retract of X , and let $i: Y \rightarrow X$ and $p: X \rightarrow Y$ be the inclusion and retraction mappings respectively. Choose a v_n self-map v of X which commutes with $i \circ p$. The map

$$p \circ v \circ i$$

is easily checked to be a v_n self-map of Y . □

COROLLARY 3.11. *The full subcategory of \mathcal{C}_1 consisting of spectra admitting a v_n self-map satisfying condition (*) of Theorem 9 is thick.*

Proof. This is similar to 3.9, and involves checking that the integers which arise in 3.6–3.8 are powers of p . In fact, the only place where an integer which is not a power of p comes up is in using 3.7 to arrange that $K(m)_*v$ is in the center of $\text{End}_{K(m)_*}(K(m)_*X)$. But this is guaranteed at the outset by condition (*). □

4. Construction of v_n self-maps

4.1. *Preliminaries.* The examples of self-maps needed for the proof of Theorem 9 are constructed using the Adams spectral sequence

$$\text{Ext}_A^{s,t}(H^*Y, H^*X) \Rightarrow [X, Y]_{t-s}$$

which relates the mod p cohomology of X and Y as modules over the Steenrod algebra to $[X, Y]_*$. The spectral sequence is usually displayed in the $(t - s, s)$ -plane, so that the groups lying in a given vertical line assemble to a single

homotopy group. With this convention the “filtration jumps” are vertical in the sense that the difference between two maps representing the same class in

$$\text{Ext}_A^{s,t}(H^*Y, H^*X)$$

represents a class in

$$\text{Ext}_A^{s',t'}(H^*Y, H^*X),$$

with $s' > s$, and $t - s = t' - s'$.

There are many criteria for convergence of the Adams spectral sequence. A simple one, which is enough for the present purpose is in [1].

LEMMA 4.1. *If X a finite spectrum and Y is a connective spectrum with the property that each $\pi_k Y$ is a finite abelian p -group, then the Adams spectral sequence converges strongly to*

$$[X, Y]_*.$$

If $B \subseteq C$ are Hopf algebras over a field k , the forgetful functor

$$C\text{-modules} \rightarrow B\text{-modules}$$

has both a left and a right adjoint. The left adjoint

$$M \mapsto C \otimes_B M$$

carries projectives to projectives, and so prolongs to a *change of rings isomorphism*

$$(4.2) \quad \text{Ext}_C^*(C \otimes_B M, N) \approx \text{Ext}_B^*(M, N).$$

When M is a C -module this can be combined with the “shearing isomorphism”

$$\begin{aligned} C \otimes_B M &\rightarrow C // B \otimes M && \left(C // B = C \otimes_B k \right), \\ c \otimes m &\mapsto \sum c'_i \otimes c''_i m, \\ \psi(c) &= \sum c'_i \otimes c''_i, \end{aligned}$$

to give another “change of rings” isomorphism

$$\text{Ext}_C^*(C // B \otimes M, N) \approx \text{Ext}_B^*(M, N).$$

The difference between Ext_C and Ext_B can therefore be measured by the augmentation ideal

$$\overline{C // B} = \ker \{ \varepsilon: C // B \rightarrow k \},$$

with the long exact sequence coming from

$$\overline{C // B} \otimes M \rightarrow C // B \otimes M \rightarrow M.$$

Recall that for $p = 2$, the dual Steenrod algebra is

$$A_* = \mathbb{F}_2[\xi_1, \xi_2, \dots], \quad |\xi_i| = 2^i - 1$$

and for p odd

$$A_* = \Lambda[\tau_0, \tau_1, \dots] \otimes \mathbb{F}_p[\xi_1, \xi_2, \dots], \quad |\tau_i| = 2p^i - 1, \quad |\xi_i| = 2(p^i - 1).$$

The subalgebra of the Steenrod algebra generated by

$$\begin{aligned} Sq^1, \dots, Sq^{2^n} & \quad \text{when } p = 2, \\ \beta, \mathcal{P}^1, \dots, \mathcal{P}^{n-1} & \quad \text{when } p \text{ is odd, and } n \neq 1, \\ \beta & \quad \text{when } p \text{ is odd and } n = 0 \end{aligned}$$

is denoted A_n . It is the finite sub-Hopf algebra which is annihilated by the ideal

$$\begin{aligned} (\xi_1^{2^{n+1}}, \xi_2^{2^n}, \dots, \xi_{n+1}, \xi_{n+2}, \dots) & \quad p = 2, \\ (\xi_1^{p^n}, \dots, \xi_n, \xi_{n+1}, \tau_{n+1}, \dots) & \quad p \neq 2. \end{aligned}$$

The augmentation ideal of $A//A_n$ is $2p^n(p - 1)$ -connected. The fact that the connectivity goes to infinity with n plays an important role in the Approximation Lemma 4.6.

It is customary to give the dual Steenrod algebra the basis of monomials in the ξ 's and τ 's. With this convention, the *Adams-Margolis elements* are

$$\begin{aligned} P_t^s & \text{ dual to } \xi_t^{p^s} \quad (s < t), \\ Q_n & \text{ dual to } \begin{cases} \tau_n & p \text{ odd} \\ \xi_{n+1} & p = 2. \end{cases} \end{aligned}$$

Each Q_n is primitive, and together they generate an exterior sub-Hopf algebra of the Steenrod algebra. The P_t^s all satisfy

$$(P_t^s)^p = 0,$$

but are primitive only when $s = 0$. The Adams-Margolis elements are naturally ordered by degree

$$\begin{aligned} |P_t^s| & = 2p^s(p^t - 1), \\ |Q_n| & = 2p^n - 1. \end{aligned}$$

4.2. *Vanishing lines.* Given an A -module M , and an Adams-Margolis element d , the *Margolis homology* of M , $H(M, d)$, is the homology of the complex (M_*, d_*) , with

$$\begin{aligned} M_n & = M, \quad n \in \mathbb{Z}, \\ d_{2n} & = d, \\ d_{2n+1} & = \begin{cases} d^{p-1} & \text{if } d = P_t^s \\ d & \text{if } d = Q_n. \end{cases} \end{aligned}$$

When X is a spectrum the symbol $H(X, d)$ will be used to denote $H(H^*X, d)$. The Margolis homology groups are periodic of period 1 if p is even or if $d = Q_n$, and they are periodic of period 2 otherwise.

Definition 4.3. Let M be an A -module. A line

$$y = mx + b$$

is a *vanishing line* of

$$\text{Ext}_A^{*,*}(M, \mathbb{F}_p)$$

if

$$\text{Ext}_A^{s,t}(M, \mathbb{F}_p) = 0 \quad \text{for } s > m(t - s) + b.$$

The following result, due to Anderson-Davis [3] and to Miller-Wilkerson [15] relates vanishing lines to Margolis homology groups.

THEOREM 4.4. *If M is a connective A -module with*

$$H(M, d) = 0 \quad \text{for } |d| \leq n,$$

then

$$\text{Ext}_A^{*,*}(M, \mathbb{F}_p)$$

has a vanishing line of slope $1/n$.

In general, there is no easy way to predict the intercept of the vanishing line, but there is the following:

PROPOSITION 4.5. *Suppose that M is a connective A -module, and that*

$$y = mx + b$$

is a vanishing line for $\text{Ext}_A^{,*}(M, \mathbb{F}_p)$. If N is a $(c - 1)$ -connected A -module, then*

$$y = m(x - c) + b$$

is a vanishing line for

$$\text{Ext}_A^{*,*}(M \otimes N, \mathbb{F}_p).$$

Proof. Let N^k be the quotient of N by the elements of degree greater than k , and $N_j^k \subseteq N^k$ the submodule consisting of elements of degree $> j$. There is an exact sequence

$$N_j^k \rightarrow N^k \rightarrow N^j.$$

Since M is connective,

$$M \otimes N = \varprojlim_k M \otimes N^k$$

and

$$\text{Ext}_A^{s,t}(M \otimes N, \mathbb{F}_p) = \varinjlim_k \text{Ext}_A^{s,t}(M \otimes N^k, \mathbb{F}_p),$$

so it suffices to prove the result for each N^k . This is trivial for $k < c$, so suppose $k \geq c$, and by induction, that the result is true for $k' < k$. Suppose that (s, t) satisfies

$$s > m(t - s - c) + b$$

and consider the exact sequence

$$M \otimes N_{k-1}^k \rightarrow M \otimes N^k \rightarrow M \otimes N^{k-1}.$$

By induction,

$$\text{Ext}_A^{s,t}(M \otimes N^{k-1}, \mathbb{F}_p) = 0.$$

The module N_{k-1}^k is just a sum of copies of $\Sigma^k \mathbb{F}_p$ —the A -module which consists of \mathbb{F}_p in degree k , and zero elsewhere. It follows that

$$\text{Ext}_A^{s,t}(M \otimes N_{k-1}^k, \mathbb{F}_p)$$

is a product of copies of

$$\text{Ext}_A^{s,t}(M \otimes \Sigma^k \mathbb{F}_p, \mathbb{F}_p) \approx \text{Ext}_A^{s,t-k}(M, \mathbb{F}_p),$$

which is zero since

$$\begin{aligned} s &> m(t - s - c) + b \\ &> m((t - k) - s) + b. \end{aligned} \quad \square$$

LEMMA 4.6 (approximation lemma). *Let M be a connective A -module, and suppose that $\text{Ext}_A^{*,*}(M, \mathbb{F}_p)$ has a vanishing line of slope m . Given b , for $n \gg 0$ the restriction map*

$$\text{Ext}_A^{s,t}(M, \mathbb{F}_p) \rightarrow \text{Ext}_{A_n}^{s,t}(M, \mathbb{F}_p)$$

is an isomorphism when

$$s \geq m(t - s) + b.$$

Proof. The result follows from the exact sequence

$$\overline{A/A_n} \otimes M \rightarrow A/A_n \otimes M \rightarrow M,$$

Proposition 4.5, and the fact that the connectivity of $\overline{A/A_n}$ can be made arbitrarily large by taking n to be large. □

4.3. *Morava K-theories and the Adams spectral sequence.* We need to be able to examine the $K(n)$ -Hurewicz homomorphism from the point of view of the Adams spectral sequence. This can be done, but it is a little easier to work with the connected cover $k(n)$ of $K(n)$. The spectrum $k(n)$ is a ring spectrum, with

$$k(n)_* = \mathbb{F}_p[v_n] \subset K(n)_* = \mathbb{F}_p[v_n, v_n^{-1}].$$

LEMMA 4.7. *The transformation $k(n)_*X \rightarrow K(n)_*X$ extends to a natural isomorphism*

$$v_n^{-1}k(n)_*X \approx K(n)_*X.$$

Proof. Since localization is exact, both sides satisfy the exactness properties of a homology theory. They agree when X is the sphere, hence for all X . □

COROLLARY 4.8. *If $k(n)_*X$ is finite then $K(n)_*X = 0$.*

Proof. If $k(n)_*X$ is finite, then for $j \gg 0$, $k(n)_jX = 0$. This means that for each $x \in k(n)_*X$, $v_n^m x = 0$ for $m \gg 0$. The result then follows from Lemma 4.7. □

Since $k(n)$ is a ring spectrum, the mod p cohomology $H^*k(n)$ is a coalgebra over the Steenrod algebra. It has been calculated by Baas and Madsen [6].

PROPOSITION 4.9. *As a coalgebra over the Steenrod algebra,*

$$H^*k(n) \approx A//E[Q_n].$$

*It follows that the E_2 -term of the Adams spectral sequence for $\pi_*k(n) \wedge X$ is isomorphic to*

$$\text{Ext}_{E[Q_n]}^{s,t}(H^*X, \mathbb{F}_p),$$

and that the map of E_2 -terms induced by the Hurewicz homomorphism is the natural restriction.

COROLLARY 4.10. *If X is a finite spectrum and $H(X, Q_n) = 0$, then $K(n)_*X = 0$.*

Proof. The group

$$\text{Ext}_{E[Q_n]}^{*,*}(H^*(X), \mathbb{F}_p)$$

is the cohomology of the complex

$$H^*X \xrightarrow{Q_n} H^*X \xrightarrow{Q_n} H^*X \xrightarrow{Q_n} \dots$$

This means that for $s > 0$, the graded abelian group

$$\text{Ext}_{E[Q_n]}^{s,*}(H^*X, \mathbb{F}_p)$$

is isomorphic to the Margolis homology group $H(X, Q_n)$. The vanishing of these groups implies that

$$\text{Ext}_{E[Q_n]}^{*,*}(H^*X, \mathbb{F}_p) \approx \text{Ext}_{E[Q_n]}^{*,0}(H^*X, \mathbb{F}_p) \subseteq H^*X$$

is finite, and hence that $k(n)_*X$ is finite. The result then follows from Corollary 4.8. □

4.4. *Examples of self maps.* The key to constructing self-maps is the following result of the second author [24]. An account appears in [21].

THEOREM 4.11. *For each $n = 1, 2, \dots$ there is a finite spectrum X_n with the properties:*

i) *All differentials in the Adams spectral sequence*

$$\text{Ext}_{E[Q_n]}^{s,t}(H^*X_n \wedge DX_n, \mathbb{F}_p) \Rightarrow k(n)_*X_n \wedge DX_n$$

are zero;

ii) *The Margolis homology groups $H(X_n \wedge DX_n, d)$ are zero if*

$$|d| < |Q_n|.$$

THEOREM 4.12. *The spectrum X_n is in $\mathcal{C}_n \setminus \mathcal{C}_{n-1}$ and has a v_n self-map satisfying (*) of Theorem 9.*

The proof of Theorem 4.12 uses the Adams spectral sequence and the following consequence of the results of Wilkerson [25].

THEOREM 4.13. *Suppose that $B \subset C$ are finite, connected, graded, co-commutative Hopf algebras over a field k of characteristic $p > 0$. If*

$$b \in \text{Ext}_B^{*,*}(k, k),$$

then for $N \gg 0$, b^{p^N} is in the image of the restriction map

$$\text{Ext}_C^{*,*}(k, k) \rightarrow \text{Ext}_B^{*,*}(k, k).$$

Proof. See Appendix A. □

Proof of Theorem 4.12. That X_n is in $\mathcal{C}_n \setminus \mathcal{C}_{n-1}$ follows from Corollary 4.10.

For the construction of the self-map, it is slightly cleaner to work from the point of view of finite ring spectra. Thus let R be the finite ring spectrum $X_n \wedge DX_n$. The ring π_*R is an algebra over π_*S^0 , and the image of π_*S^0 in

π_*R is in the center (in the graded sense). Similarly, if $B \subseteq A$ is a sub-Hopf algebra, the ring

$$\text{Ext}_B^{*,*}(H^*R, \mathbb{F}_p)$$

is a central algebra over $\text{Ext}_B^{*,*}(\mathbb{F}_p, \mathbb{F}_p)$.

To show that X_n admits a v_n self-map satisfying condition (*) of Theorem 9 it suffices to exhibit an element

$$v \in \pi_*R$$

satisfying

$$(4.14) \quad k(n)_*v^{p^M} = v_n^{p^N} \cdot 1, \quad \text{for some } M, N > 0;$$

$$(4.15) \quad \text{the map } k(m)_*v \text{ is nilpotent when } m \neq n.$$

Step 1. The first step is to find an approximation to a v_n self-map in the E_2 -term of the Adams spectral sequence. Let

$$v_n \in \text{Ext}_{E[Q_n]}^{1, 2p^n - 1}(\mathbb{F}_p, \mathbb{F}_p)$$

be the element representing $v_n \in k(n)_*$. We need to find a

$$w \in \text{Ext}_A^{p^N, p^N(2p^n - 1)}(H^*R, \mathbb{F}_p)$$

restricting to $v_n^{p^N} \cdot 1$, for $N \gg 0$. By 4.4, the bigraded group

$$\text{Ext}_A^{*,*}(H^*R, \mathbb{F}_p)$$

has a vanishing line of slope $1/2(p^n - 1)$. Using the approximation lemma, an integer n can be chosen for which the restriction map

$$(4.16) \quad \text{Ext}_A^{s,t}(H^*R, \mathbb{F}_p) \rightarrow \text{Ext}_{A_n}^{s,t}(H^*R, \mathbb{F}_p)$$

is an isomorphism if

$$s > \frac{1}{2(p^n - 1)}(t - s).$$

By Theorem 4.13 there is an element

$$\tilde{w} \in \text{Ext}_{A_n}^{*,*}(\mathbb{F}_p, \mathbb{F}_p)$$

restricting to $v_n^{p^N} \in \text{Ext}_{E[Q_n]}^{*,*}(\mathbb{F}_p, \mathbb{F}_p)$. The class w can be taken to be the image of $\tilde{w} \cdot 1$ under the isomorphism (4.16). \square

Step 2. This construction of the class w actually gives something more. Since

$$\text{Ext}_{A_n}^{*,*}(\mathbb{F}_p, \mathbb{F}_p)$$

is in the center (in the graded sense) of

$$\text{Ext}_{A_n}^{*,*}(H^*R, \mathbb{F}_p),$$

the class w commutes with every

$$\alpha \in \text{Ext}_A^{s,t}(H^*R, \mathbb{F}_p)$$

with

$$(4.17) \quad s \geq \frac{1}{2(p^n - 1)}(t - s).$$

Step 3. Now we choose a power of w which survives the Adams spectral sequence. The differentials in the Adams spectral sequence are derivations, and the values of $d_r w$ lie in the region (4.17). This means that

$$d_{r-1}w = 0 \Rightarrow d_r w^p = 0.$$

Since $d_1 w = 0$ it follows that $d_b w^{p^b} = 0$. The possible values of $d_r w^{p^b}$ for $r > b$ lie in the region

$$s > \frac{1}{2(p^n - 1)}(t - s),$$

which is above the vanishing line. This means that the class w^{p^b} is a permanent cycle.

Step 4. For simplicity, replace w with w^{p^b} , and adjust the integer N so that w restricts to $v_n^{p^N} \cdot 1$. Let

$$v \in \pi_* R$$

be a representative of w . We will see that this is the desired class.

The difference $k(n)_*(v - v_n^{p^N})$ is represented by a class in

$$\text{Ext}_{E[Q_n]}^{s,t}(H^*R, \mathbb{F}_p)$$

with $s > 1/2(p^n - 1)(t - s)$. Some power of $k(n)_*(v - v_n^{p^N})$ is therefore represented by a class above the vanishing line of

$$\text{Ext}_{E[Q_n]}^{*,*}(H^*R, \mathbb{F}_p)$$

(which has slope $1/2(p^n - 1)$), and hence is zero. Lemma 3.4 then gives that

$$k(n)_* v^{p^M} = v_n^{p^{MN}}, \quad M \gg 0.$$

This proves property (4.14).

Property (*) is trivial when $m < n$, since $R \in \mathcal{C}_n$. When $m > n$, it is a consequence of Lemma 2.1 and the fact that the Adams spectral sequence for $k(m)_* R$ has a vanishing line of slope $1/2(p^m - 1)$, and that the powers of v are represented by classes lying on the line

$$s = \frac{1}{2(p^n - 1)}(t - s)$$

which has a larger slope. This completes the proof. □

4.5. *Proof that Theorem 7 implies Theorem 2.* This subsection is included to satiate any curiosity aroused by the claim made after the statement of Theorem 7. Since the argument is not necessary for establishing any of the results of this paper, it is included only as a sketch.

In [11, §1] the nilpotence theorem (Theorem 2 in this paper) is reduced to showing that if R is a connective, associative ring spectrum, and

$$\alpha \in \pi_* R$$

is in the kernel of the MU -Hurewicz homomorphism, then α is nilpotent. This in turn is easily reduced to the case when R is localized at p and MU is replaced with BP . The case $|\alpha| \leq 0$ is easy, so it may be assumed that $|\alpha| > 0$.

Let $\bar{\alpha}$ be the map

$$(4.18) \quad \Sigma^{|\alpha|} R \xrightarrow{\alpha \wedge 1} R \wedge R \rightarrow R.$$

The map (4.18) induces multiplication by $BP_* \alpha = 0$ in BP homology. This means that from the point of view of the Adams-Novikov spectral sequence, composition with $\bar{\alpha}$ moves the homotopy groups to the right along a line of positive slope.

Step 1. The construction used to produce the spectra X_n of this section can be used to construct finite torsion-free spectra Y_n with the property that $H^* Y_n$, as a module over A_n , is free over $A_n // E$, where E is the sub-Hopf algebra

$$\Lambda[Q_0, \dots, Q_n].$$

See [24].

Step 2. Use the spectral sequence of [20, Th. 4.4.3] to show that

$$\text{Ext}_{BP_* BP}^{s,t}(BP_*, BP_* R \wedge Y_n)$$

has a vanishing line with slope tending to zero as $n \rightarrow \infty$.

Step 3. It follows from the vanishing line that for $n \gg 0$ the spectrum

$$Y_n \wedge \alpha^{-1} R$$

is contractible.

Step 4. Now use Theorem 14 to conclude that Y_n is Bousfield equivalent to the sphere, hence that $\alpha^{-1} R$ is contractible, hence that α is nilpotent.

5. Endomorphisms, up to nilpotents

In this section we will completely classify the endomorphisms (up to nilpotent elements) of full subcategories of finite spectra which are stable under suspension (Theorem 5.3). This has as a consequence Theorem 11, which is essentially the special case in which the subcategory consists of one object.

5.1. *N-endomorphisms.* The v_n self-maps form an endomorphism (up to nilpotent elements) of the category \mathcal{C}_n . It turns out that these are the only endomorphisms of this kind that can occur in the category of finite spectra.

Definition 5.1. Let \mathcal{C} be a full subcategory of \mathcal{C}_0 which is closed under suspensions. A collection v , of self-maps

$$v_X: \Sigma^{k_X} X \rightarrow X, \quad X \in \mathcal{C}$$

is said to *represent an N-endomorphism of C* if

i) The map $v_{\Sigma X}$ is the composite

$$\begin{array}{ccc} \Sigma^k \Sigma X & \xrightarrow{\text{flip}} & \Sigma^k X \wedge S^1 \\ & & \downarrow v_X \wedge 1_{S^1} \\ & & X \wedge S^1 \xrightarrow{\text{flip}} \Sigma X. \end{array}$$

ii) For each $f: X \rightarrow Y$ in \mathcal{C} there are integers i and j with $ik_X = jk_Y$, such that the following diagram commutes:

$$\begin{array}{ccc} \Sigma^M X & \xrightarrow{f} & \Sigma^M Y \\ v_X^i \downarrow & & \downarrow v_Y^j \\ X & \xrightarrow{f} & Y. \end{array}$$

Two representatives v and v' are *equivalent* if for each $X \in \mathcal{C}$ there are integers i and j with $v_X^i = v_X'^j$. An *N-endomorphism* is defined to be an equivalence class of representatives. The *N-endomorphism* represented by v will be denoted $[v]$.

Remark 1. (1) If v represents an *N-endomorphism* of a category \mathcal{C} , and $f: X \rightarrow Y$ an isomorphism with $X \in \mathcal{C}$, then defining v_Y to be

$$\begin{array}{ccc} \Sigma^k Y & \xrightarrow{\Sigma^k f^{-1}} & \Sigma^k X \\ & & \downarrow v_X \\ & & X \xrightarrow{f} Y \end{array}$$

extends v to the full subcategory obtained from \mathcal{C} by adjoining the suspensions of Y . Because of property ii), the resulting *N-endomorphism* is independent of the choice of isomorphism f . In this way an *N-endomorphism* can always be extended uniquely to a full subcategory which is closed under suspensions

and isomorphisms. This procedure will be used without comment, so once an N -endomorphism has been defined on a subcategory C of finite spectra, it will be taken to be extended to the smallest full subcategory containing C , which is closed under suspensions and isomorphisms. Among other things, this means that if v_X is defined, so is $v_{X \wedge S^1}$ and, after raising both sides to a power if necessary, one has

$$v_{X \wedge S^1} = v_X \wedge 1_{S^1}.$$

(2) An N -endomorphism is of *degree zero* if all of the integers k_X are zero. If an N -endomorphism is not of degree zero, then none of the integers k_X is zero, and the maps v_X can all be chosen to have finite order. Given two spectra $X, Y \in C$, the maps v_X and v_Y can be chosen in such a way that the integers k_X and k_Y coincide. With this arrangement, given a map

$$f \in [X, Y]_*$$

if there are integers i and j for which

$$v_Y^i \circ f = f \circ v_X^j,$$

then it must be the case that $i = j$. This same discussion applies to any finite collection of elements of C .

Example 5.2. (1) Any collection v with each v_X nilpotent represents an N -endomorphism.

(2) Taking each v_X to be a multiple of the identity defines an actual endomorphism.

(3) Suppose $C \subseteq C_n$. Taking v_X to be a v_n self-map defines an N -endomorphism.

5.2. *Classification of N -endomorphisms.* The above list of examples turns out to be complete.

THEOREM 5.3. *Let $C \subseteq C_0$ be a full subcategory which is closed under suspensions. If v represents an N -endomorphism of C then one of the following holds:*

- i) *each v_X is nilpotent;*
- ii) *some power of each v_X is a multiple of the identity map;*
- iii) *$C \subseteq C_n$ for some n , and each v_X is a v_n self-map.*

Of course, these possibilities are not mutually exclusive. If $X \in C_{n+1} \subseteq C_n$, any v_n self-map of X is nilpotent.

The proof of Theorem 5.3 will be completed in subsection 5.6..

COROLLARY 5.4. *Suppose that $X \in \mathcal{C}_0$, and that $v \in [X, X]_*$ is in the center. Then v is nilpotent, a power of v is a multiple of the identity, or v is a v_n self-map.*

Proof. Let \mathcal{C} be the full subcategory of \mathcal{C}_0 consisting of the suspensions of X . The map v represents an N -endomorphism of \mathcal{C} , so the result follows from Theorem 5.3. □

Proof of Theorem 11. Let $X \in \mathcal{C}_n \setminus \mathcal{C}_{n+1}$. By Corollary 5.4 the map induced by the $K(n)$ -Hurewicz homomorphism

$$\text{center } [X, X]_* \approx_F \begin{cases} \mathbb{Z}_{(p)} & (n = 0) \\ \mathbf{F}_p[v_n] & (n \neq 0) \end{cases}$$

has a nilpotent kernel. The result then follows from Theorem 9 and the observation made in the proof of Corollary 3.11 that some p^N power of any v_n self-map satisfying condition (*) of Theorem 9 is central. □

Remark 2. It is natural to consider a representative v of an N -endomorphism for which the integers i and j can be taken to be powers of p , and to define a finer equivalence relation by declaring v and v' to be equivalent if for each $X \in \mathcal{C}$ there are integers i and j with $v_X^{p^i} = v'_X{}^{p^j}$. An F -endomorphism is an equivalence class of such representatives with respect to this finer equivalence relation. Taking v_X to be a v_n self-map satisfying condition (*) of Theorem 9 defines an F -endomorphism.

The argument which classifies N -endomorphisms leads to a classification of F -endomorphisms, provided one keeps track of when certain integers can be taken to be a power of p . We have kept track of this in the statements of the results leading to the proof of Theorem 5.3, but since we have no use for F -endomorphisms in mind, we leave the classification to the interested reader.

The proof of Theorem 5.3 falls into two parts. First it is shown that an N -endomorphism extends uniquely to a thick subcategory. It then suffices to construct, for each n , a spectrum $X_n \in \mathcal{C}_n \setminus \mathcal{C}_{n+1}$ whose only non-nilpotent self-map is a v_n self-map.

To begin, we dispense with the N -endomorphisms of degree zero.

PROPOSITION 5.5. *If X is in \mathcal{C}_0 , and $v: X \rightarrow X$ is in the center of $[X, X]_* = \pi_* X \wedge DX$ then there are integers m and n for which*

$$v^n = \text{multiplication by } m.$$

Proof. Since

$$\pi_* X \wedge DX \otimes \mathbb{Q} \approx H\mathbb{Q}_* X \wedge DX \approx \text{End } H\mathbb{Q}_* X,$$

the map $H\mathbb{Q}_*v$ must be in the center of $\text{End } H\mathbb{Q}_*X$ which consists entirely of scalar multiplications. Since the Hurewicz map $H\mathbb{Q}_*$ factors through $H\mathbb{Z}_*$, this constant must be an integer k . The map $w = v - k$ then has finite order.

Since all of the eigenvalues of $H\mathbb{F}_{p^*}w^{p-1}$ are equal to 0 or 1, the map $H\mathbb{F}_{p^*}w^{(p-1)p^M}$ is an idempotent for $M \gg 0$. Replace w with $w^{(p-1)p^M}$. The map w still has finite order and is in the center of $[X, X]_*$. Define connective spectra A_1 and A_2 by

$$A_1 = w^{-1}X, \quad A_2 = (1 - w)^{-1}X.$$

The map

$$X \rightarrow A_1 \vee A_2$$

induces an isomorphism on both mod p and rational homology, hence on homology with coefficients in $\mathbb{Z}_{(p)}$. It is therefore a homotopy equivalence, and in particular A_1 and A_2 are finite.

The ring of self-maps $[X, X]_*$ can be written as a ring of 2×2 matrices, in which the ij -entry is in $[A_j, A_i]$. The map w is represented by the matrix

$$\begin{pmatrix} w|_{A_1} & 0 \\ 0 & 0 \end{pmatrix},$$

whose $(1, 1)$ entry is an equivalence. Given a map $f: \Sigma^k A_2 \rightarrow A_1$, let \tilde{f} be the map

$$\begin{pmatrix} 0 & f \\ 0 & 0 \end{pmatrix}.$$

Then

$$\text{ad}(w)\tilde{f} = \begin{pmatrix} w|_{A_1}f & 0 \\ 0 & 0 \end{pmatrix}.$$

Since w is central, and $w|_{A_1}$ is an equivalence this means that f is null. By Lemma 5.6 below, it follows that one of A_1 and A_2 is contractible. If A_1 is contractible, then w is nilpotent, and the result follows from Lemma 3.4. If A_2 is contractible, then $1 - w$ is nilpotent, $H\mathbb{Q}_*X = 0$, and we may assume that the integer k is 0, so that $w = v^{(p-1)p^M}$. It then follows from Lemma 3.4 that $v^{(p-1)p^M} = 1$ for $M \gg 0$. This completes the proof. \square

We have used

LEMMA 5.6. *If A and B are non-contractible p -local finite spectra, then $[A, B]_* \neq 0$.*

Proof. Since

$$H\mathbb{F}_{p^*}DA \wedge B = \text{Hom}(H\mathbb{F}_{p^*}A, H\mathbb{F}_{p^*}B) \neq 0,$$

the spectrum $DA \wedge B$ is not-contractible. It therefore has a nonzero homotopy group. Now use the isomorphism

$$\pi_* DA \wedge B \approx [A, B]_* \quad \square$$

5.3. *Some technical tools.* The next few results are a bit technical, but they come up several times.

LEMMA 5.7. *Suppose that M is a bimodule over the ring $\mathbb{Z}_{(p)}[v]$, and for $m \in M$ let*

$$\text{ad}(v)m = vm - mv.$$

If there are integers i, j and k , for which

i) $k \text{ad}(v^i)m = 0$, and

ii) $\text{ad}(v^j)(\text{ad}(v^i)m) = 0$,

then

$$\text{ad}(v^{ijk})m = 0.$$

Clearly, k can be taken to be a power of p , so that if i and j are powers of p , then so is ijk .

LEMMA 5.8. *Suppose M is a bimodule over the ring $\mathbb{Z}[v]$. Let $\text{ad}(v): M \rightarrow M$ be the operator $\text{ad}(v)m = vm - mv$. The operators $\text{ad}(v^n)$ and $\text{ad}^i(v)$ are related by the formula:*

$$(5.9) \quad \text{ad}(v^n)m = \sum_{i < n} \binom{n}{i} \text{ad}^i(v)m \cdot v^{n-i}.$$

Proof. Let $l(v)$ and $r(v)$ be the operators of left and right multiplication by v respectively. Then

$$\text{ad}(v) = l(v) - r(v),$$

and the operators $\text{ad}(v)$, $l(v)$, and $r(v)$ all commute. Now take the equation

$$l(v) = \text{ad}(v) + r(v),$$

raise both sides to the power n , and use the binomial theorem. □

Proof of 5.7. Replacing j with ij and v with v^i , we may suppose that $i = 1$. Since the operators $\text{ad}(v^j)$ and $\text{ad}(v)$ commute, it follows from (5.9) that

$$\text{ad}(v^j)(\text{ad}(v^j)m) = 0.$$

Replacing v with v^j , we may therefore also assume that $j = 1$, and hence that $\text{ad}^2(v)m = 0$. But then, again from (5.9),

$$\text{ad}(v^k)m = k \text{ad}(v)m = 0.$$

This completes the proof. □

5.4. *N*-endomorphisms and thick subcategories.

LEMMA 5.10. *Suppose that $[v]$ and $[v']$ are N -endomorphisms of a thick subcategory $\mathcal{C} \subseteq \mathcal{C}_0$. The full subcategory $\mathcal{D} \subseteq \mathcal{C}$ consisting of objects for which $[v] = [v']$ is thick.*

Proof. The case in which the degree of $[v]$ is zero follows from Proposition 5.5, so we may assume that the degree of $[v]$ is not zero. In this case Remark 2 applies.

Suppose that $i: Y \rightarrow X$ is the inclusion of a retract, with $X \in \mathcal{D}$ and $Y \in \mathcal{C}$. Choose the representatives v_X, v_Y, v'_X and v'_Y , so that

$$(1) \ v_X i = i v_Y, \ v'_X i = i v'_Y, \text{ and}$$

$$(2) \ v_X = v'_X.$$

Then $i v_Y = i v'_Y$, so that $v_Y = v'_Y$ since i is a monomorphism.

Now suppose that

$$X \rightarrow Y \rightarrow Z$$

is a cofiber sequence in \mathcal{C} with X and Y in \mathcal{D} . Choose the maps v to have finite order, with $v_X = v'_X, v_Y = v'_Y$, and so that

$$\begin{array}{ccccc} \Sigma^M Y & \longrightarrow & \Sigma^M Z & \xrightarrow{\delta} & \Sigma^M X \wedge S^1 \\ v_Y \downarrow & & v_Z \downarrow & & v_{X \wedge S^1} \downarrow \\ Y & \longrightarrow & Z & \xrightarrow{\delta} & X \wedge S^1 \end{array}$$

and

$$\begin{array}{ccccc} \Sigma^M Y & \longrightarrow & \Sigma^M Z & \xrightarrow{\delta} & \Sigma^M X \wedge S^1 \\ v'_Y \downarrow & & v'_Z \downarrow & & v'_{X \wedge S^1} \downarrow \\ Y & \longrightarrow & Z & \xrightarrow{\delta} & X \wedge S^1. \end{array}$$

commute. Make $[Z, Z]_*$ into a $\mathbb{Z}_{(p)}[v]$ -bimodule by

$$(5.11) \quad v m = v_Z \circ m,$$

$$(5.12) \quad m v = m \circ v'_Z.$$

The goal is to show that $\text{ad}(v^k)1_Z$ is zero, for some k . We know that $\text{ad}(v^k)1_Z$ has finite order, and that it factors through a map $\Sigma^M X \wedge S^1 \rightarrow Z$. In general, if a map $f \in [Z, Z]_*$ factors through some $W \in \mathcal{D}$ then $\text{ad}(v^j)f = 0$ for some j . It follows that

$$\text{ad}(v^j)(\text{ad}(v)1_Z) = 0$$

for some j . The result then follows from Lemma 5.7. □

LEMMA 5.13. *Let $[v]$ be an N -endomorphism of a full subcategory \mathcal{C} of \mathcal{C}_0 which is closed under suspensions. If \mathcal{D} is the largest full subcategory of \mathcal{C}_0 to which v extends, then \mathcal{D} is thick.*

Proof. If the degree of $[v]$ is zero, then the result follows from 5.5. We may therefore assume that the maps v all have finite order, and for the finitely many spectra that come up in the proof, that they all have the same degree. Suppose that $X \in \mathcal{D}$ and that

$$\begin{aligned} i: Y &\rightarrow X, \\ p: X &\rightarrow Y \end{aligned}$$

satisfy $p \circ i = 1_Y$. Choose the map v_X so that it commutes with the idempotent $i \circ p$, and set

$$v_Y = p \circ v_X \circ i.$$

If it can be shown that this map v_Y extends $[v]$ to the full subcategory $\mathcal{D} \cup \{\Sigma^k Y\}_{k \in \mathbb{Z}}$, it will follow that $Y \in \mathcal{D}$ by maximality.

To check this, let $W \in \mathcal{D} \cup \{\Sigma^k Y\}_{k \in \mathbb{Z}}$ and suppose at first that $W \in \mathcal{D}$. Given a map $f: W \rightarrow Y$, choose an integer j so that the outer rectangle in the following diagram commutes:

$$\begin{array}{ccccc} \Sigma^N W & \xrightarrow{f} & \Sigma^N Y & \xrightarrow{i} & \Sigma^N X \\ v_W^j \downarrow & & v_Y^j \downarrow & & v_X^j \downarrow \\ W & \xrightarrow{f} & Y & \xrightarrow{i} & X. \end{array}$$

The right square commutes by construction, so the left must also, since the map i is the inclusion of a wedge summand. The argument for dealing with a map $Y \rightarrow W$ is similar. Finally, given

$$f: \Sigma^k Y \rightarrow Y,$$

using the above, find an integer k so that

$$i \circ v_Y^k \circ f = v_X^k \circ i \circ f = i \circ f \circ v_Y^k.$$

Then, again, since $i_*: [Y, Y]_* \rightarrow [Y, X]_*$ is a monomorphism,

$$v_Y^k \circ f = f \circ v_Y^k.$$

Now suppose that

$$X \rightarrow Y \rightarrow Z$$

is a cofiber sequence with X and Y in \mathcal{D} . Choose the maps v_X and v_Y so that the left square in the following diagram commutes:

$$\begin{array}{ccccc} \Sigma^N X & \longrightarrow & \Sigma^N Y & \longrightarrow & \Sigma^N Z \\ v_X \downarrow & & v_Y \downarrow & & v_Z \downarrow \\ X & \longrightarrow & Y & \longrightarrow & Z, \end{array}$$

Let v_Z be any map of finite order making the above a map of cofiber sequences. If it can be shown that this map v_Z extends v to the category

$$\mathcal{D} \cup \{\Sigma^k Z\}_{k \in \mathbb{Z}},$$

it will follow that $Z \in \mathcal{D}$ by maximality.

To check this, suppose that $f: Z \rightarrow W$ is a map with W an object of $\mathcal{D} \cup \{\Sigma^k Z\}_{k \in \mathbb{Z}}$. If necessary raise the maps v_W, v_X, v_Y , and v_Z to powers so that they are all of finite order, and that they all have the same degree.

Case 1. $W \in \mathcal{D}$, and f factors through $\delta: Z \rightarrow X \wedge S^1$. Write

$$f = g \circ \delta,$$

and choose an integer i so that the right square in the following diagram commutes:

$$\begin{array}{ccccc} \Sigma^N Z & \xrightarrow{\delta} & \Sigma^N X \wedge S^1 & \xrightarrow{g} & \Sigma^N W \\ v_Z^i \downarrow & & v_X^i \downarrow & & \downarrow v_W^i \\ Z & \xrightarrow{\delta} & \Sigma X & \xrightarrow{g} & W. \end{array}$$

The left square commutes by definition of v_Z , so that the whole diagram commutes. □

Case 2. *The spectrum W is in \mathcal{D} .* Make the graded abelian group $[Z, W]_*$ into a bimodule over $\mathbb{Z}_{(p)}[v]$ by

$$\begin{aligned} v \cdot m &= v_W \circ m, \\ m \cdot v &= m \circ v_Z. \end{aligned}$$

For some i , $\text{ad}(v^i)f$ vanishes on Y and so factors through $X \wedge S^1$. By Case 1,

$$\text{ad}(v^j) (\text{ad}(v^i)f) = 0 \quad \text{for some } j,$$

and the result follows from Lemma 5.7.

Case 3. $W = \Sigma^k Z$. By Case 2, the maps v can be chosen to commute in addition with all elements of $[Z, \Sigma^k X \wedge S^1]$. It then follows that $\text{ad}(v)f$ factors through $\Sigma^k Y$, and so by case 1,

$$\text{ad}(v^j) (\text{ad}(v)f) = 0 \quad \text{for some } j.$$

The result then follows from Lemma 5.7.

The case of maps $W \rightarrow Z$ is handled similarly. This completes the proof. □

5.5. *A spectrum with few non-nilpotent self maps.* Now to construct, for each n a spectrum $X_n \in \mathcal{C}_n \setminus \mathcal{C}_{n-1}$ whose only non-nilpotent self-maps are roots of the identity, or v_n self-maps. The spectrum X_0 can be taken to be the sphere.

PROPOSITION 5.14. *For each $n > 0$ there exists a sequence*

$$\underline{k} = (k_0, \dots, k_{n-1}),$$

and a finite spectrum $M(\underline{k}) \in \mathcal{C}_n \setminus \mathcal{C}_{n-1}$, satisfying:

- i) $BP_*M(\underline{k}) = BP_*/(v_0^{p^{k_0}}, \dots, v_{n-1}^{p^{k_{n-1}}}) \quad (v_0 = p)$;
- ii) If

$$v: \Sigma^j M(\underline{k}) \rightarrow M(\underline{k})$$

is a non-nilpotent self-map, then some power of v is the identity map, or v is a v_n self-map.

Proof. Suppose by induction on n that a sequence

$$\underline{k} = (k_0, \dots, k_{n-1})$$

and a spectrum

$$M = M(\underline{k})$$

have been found, satisfying condition i). When $n = 1$ the sequence can be taken to be (1), and the spectrum M , the mod p Moore spectrum

$$S^0 \cup_p e^1.$$

Let $I(\underline{k}) \subset BP_*$ be the ideal

$$(v_0^{p^{k_0}}, \dots, v_{n-1}^{p^{k_{n-1}}}).$$

If v is a non-nilpotent self-map of $M(\underline{k})$ then the BP -Hurewicz image, BP_*v , must be a non-nilpotent element of the ring

$$\text{Hom}_{BP_*}(BP_*/I(\underline{k}), BP_*/I(\underline{k})) \approx BP_*/I(\underline{k}).$$

The map BP_*v must also be a map of BP_*BP -comodules, and so is an element of

$$\text{Hom}_{BP_*BP}(BP_*/I(\underline{k}), BP_*/I(\underline{k})) \subset BP_*/I(\underline{k}).$$

Now the ideal

$$I_n = (p, v_1, \dots, v_{n-1}) \subset BP_*/I(\underline{k})$$

is nilpotent, and one knows ([20, Th. 4.3.2]) that

$$\text{Hom}_{BP_*BP}(BP_*/I_n, BP_*/I_n) = \mathbb{F}_p[v_n].$$

It follows that there are elements $\lambda \in \mathbb{F}_p$, $k \in \mathbb{Z}$, with the property that $BP_*v - \lambda v_n^k$ is an element of I_n , hence nilpotent. It then follows from Lemma 3.4 that

$$BP_*v^{(p-1)p^N} = v_n^{k(p-1)p^N}, \quad N \gg 0.$$

Replace v with $v^{(p-1)p^N}$. If $k = 0$, then $BP_*(v-1) = 0$, and so $v-1$ is nilpotent (by Theorem 2). It then follows from Lemma 3.4 that

$$v^{p^N} = 1_M, \quad N \gg 0.$$

Suppose then that $k \neq 0$, and let w be a v_n self-map of M . By the above discussion applied to w , there are integers i and j , for which $BP_*v^i = BP_*w^j$. But this means (again by Theorem 2) that $v^i - w^j$ is nilpotent, so by Lemma 3.4 some power of v is homotopic to some power of w , and v is a v_n self-map. This proves iii). For the rest of the induction step, let

$$w: \Sigma^{2(p^n-1)p^N} M \rightarrow M$$

be a v_n self-map satisfying condition (*) of Theorem 9. The integer k_n can be taken to be N , and

$$M(k_0, \dots, k_n),$$

the cofiber of the map w . □

5.6. *Proof of Theorem 5.3.* Let v be an N -endomorphism of $\mathcal{C} \subseteq \mathcal{C}_0$. Then v extends uniquely to the smallest thick subcategory $\mathcal{C}_n \subset \mathcal{C}_0$ containing \mathcal{C} . Let

$$\underline{k} = (k_0, \dots, k_{n-1})$$

and $M = M(\underline{k})$ be as in Lemma 5.14, and let \mathcal{D} be the full subcategory of \mathcal{C}_n consisting of the suspensions of M . Then v is also uniquely determined by its restriction to \mathcal{D} , i.e. by the map v_M . By Proposition 5.14, there are three possibilities for v_M , and these are the restrictions of the nilpotent, identity, and v_n self-map N -endomorphisms. This completes the proof of Theorem 5.3. □

Appendix A. Proof of Theorem 4.13

The purpose of this appendix is to prove (rather, deduce from [25]) Theorem 4.13. All of the techniques used here can be found in [25].

Throughout this appendix, all Hopf algebras will be over a field of characteristic $p > 0$. They will be connected, graded, cocommutative, and finite-dimensional. The dual of a Hopf algebra will be graded in such a way that the dual of the homogeneous component of degree k has degree $-k$. This convention enables the coaction map (A.2) to preserve degrees.

The action of a Hopf algebra B on a module M can be expressed as an “action”

$$(A.1) \quad B \otimes M \rightarrow M$$

or as a “coaction”

$$(A.2) \quad M \rightarrow B^* \otimes M.$$

A module M which happens to be an algebra is an *algebra over B* if the multiplication map

$$M \otimes M \rightarrow M$$

is a map of B -modules. This is equivalent to the requirement that the coaction map (A.2) be multiplicative. All algebras over Hopf algebras in this appendix will be graded and connected.

If $B \subseteq C$ is normal, and M is a C -module, then the sub-module of elements invariant under B ,

$$M^B = \text{Hom}_B(k, M),$$

inherits an action of the quotient Hopf algebra $C//B$. In fact, so do all of the derived functors

$$(A.3) \quad \text{Ext}_B^*(k, M).$$

If M is an algebra over B , then (A.3) becomes an algebra over $C//B$ [23].

In case $B \subseteq C$ is normal, the relationship between the cohomologies of B and C is given by the Lyndon-Hochschild-Serre spectral sequence

$$\text{Ext}_{C//B}^*(k, \text{Ext}_B^*(k, M)) \Rightarrow \text{Ext}_C^*(k, M).$$

The main result of this appendix is

THEOREM A.4. *Suppose that R is a Noetherian C -algebra, and that $B \subseteq C$ is normal. Then*

- i) *The algebra $\text{Ext}_C^*(k, R)$ is Noetherian, hence finitely generated.*
- ii) *The Lyndon-Hochschild-Serre spectral sequence*

$$\text{Ext}_{C//B}^*(k, \text{Ext}_B^*(k, R)) \Rightarrow \text{Ext}_C^*(k, R)$$

terminates at a finite stage in the sense that there is an integer N with the property that all of the differentials d_r are zero, if $r > N$.

- iii) *There is an integer N with the property that $d_r x^{p^N}$ is zero, for all x and all r .*

- iv) *The Lyndon-Hochschild-Serre spectral sequence is of finitely generated modules over some connected, graded, Noetherian ring T .*

The parts of this theorem are closely related.

LEMMA A.5. *In Theorem A.4, parts i), ii), and iii) follow from iv). Given i), parts ii), iii), and iv) are equivalent.*

Proof. Suppose first that iv) holds. Then part ii) follows from Lemma A.6 below. Part iii) follows from ii) since the differentials are derivations. That iv) \Rightarrow i) follows from the fact that if a ring is complete with respect to an exhaustive filtration, and if the associated graded ring is Noetherian, then so is the original ring (see [8, 3.2.9 and Cor. 2 to Prop. 12] or [4, Cor. 10.25]).

Now suppose that part i) holds. Then the the E_2 -term of the Lyndon-Hochschild-Serre spectral sequence is Noetherian, hence finitely generated over k . Given ii), part iii) follows as above. Given part iii), the algebra T in part iv) can be taken to be the algebra of $(p^N)^{\text{th}}$ -powers in E_2 . The implication iv) \Rightarrow ii) was established in the preceding paragraph. This completes the proof. \square

We have used:

LEMMA A.6. *Let $\{E_r, d_r\}$ be a spectral sequence of finitely generated modules over a Noetherian ring T . There is an integer N with the property that all of the differentials d_r are zero, if $r > N$.*

Proof. The modules E_r are sub-quotients of E_2 . Define

$$B_{r+1} \subseteq B_r \subseteq \cdots \subseteq Z_r \subseteq Z_{r+1} \cdots \subseteq E_2$$

with the property that

$$E_{r+1} = Z_r/B_r.$$

The graded T -modules Z_r and B_r can be thought of as the kernel and image of d_r respectively. By the ascending chain condition, there is an integer N with the property that $B_r = B_N$ if $r \geq N$. But this implies, for $r \geq N + 1$ that $E_r \subseteq E_{N+1}$, so that the image of d_r is zero. \square

Wilkerson [25] has proved a special case of Theorem A.4. Recall that a map $R \rightarrow S$ of rings is finite if S is finitely generated when regarded as an R -module.

THEOREM A.7 (Wilkerson). *i) Suppose that $B \subseteq C$ is in the center, and that the action of C on R is trivial. Then i)–iv) of Theorem A.4 hold.*

ii) If $B \subseteq C$ is normal, the map

$$\text{Ext}_C^*(k, k) \rightarrow \text{Ext}_B^*(k, k)$$

is finite.

The requirement that C act trivially on R turns out not to be much of a restriction.

LEMMA A.8. *Suppose a Hopf algebra A acts on a graded commutative ring R . Given an element $x \in R$, for $N \gg 0$, the element x^{p^N} is invariant under A . In particular, if R is Noetherian, then*

$$R^A \hookrightarrow R$$

is finite.

Proof. This is easiest to verify from the point of view of the coaction. By assumption, the coaction is given by

$$\psi(x) = 1 \otimes x + \sum a_i \otimes x_i,$$

where $|x_i| \neq 0$. Since A is finite-dimensional, there is an N with the property that $a^{p^N} = 0$ for all $a \in A^*$ with $|a| \neq 0$. But then

$$\begin{aligned} \psi(x^{p^N}) &= 1 \otimes x^{p^N} + \sum a_i^{p^N} \otimes x_i^{p^N} \\ &= 1 \otimes x^{p^N}. \end{aligned}$$

This completes the proof. □

LEMMA A.9. *If A is a finite Hopf algebra and $R \rightarrow S$ is a finite map of Noetherian A -algebras, then*

$$\text{Ext}_A^*(k, R) \rightarrow \text{Ext}_A^*(k, S)$$

is finite.

COROLLARY A.10. *If C is a Hopf algebra, and R is a Noetherian C -module, then the cohomology algebra*

$$\text{Ext}_C^*(k, R)$$

is Noetherian.

Proof. By Lemma A.8, the map $R^C \rightarrow R$ is finite. By Lemma A.9,

$$R^C \otimes \text{Ext}_C^*(k, k) \approx \text{Ext}_C^*(k, R^C) \rightarrow \text{Ext}_C^*(k, R)$$

is finite. The result now follows from A.7. □

COROLLARY A.11. *It suffices to prove Theorem A.4 when $R = k$.*

Proof. It is enough to deduce part iv) of Lemma A.4 since by Lemma A.5, iv) implies i), ii), and iii). Suppose that the Lyndon-Hochschild-Serre spectral sequence

$$\text{Ext}_{C//B}^*(k, \text{Ext}_B^*(k, k)) \Rightarrow \text{Ext}_C^*(k, k)$$

consists of finitely generated modules over the Noetherian ring T . Then the spectral sequence

$$\text{Ext}_{C//B}^*(k, \text{Ext}_B^*(k, R^C)) \Rightarrow \text{Ext}_C^*(k, R^C)$$

consists of finitely generated modules over the Noetherian ring $R^C \otimes T$. By Lemma A.8, the map $R^C \rightarrow R$ is finite. It follows from Lemma A.9 that the map

$$\text{Ext}_{C//B}^*(k, \text{Ext}_B^*(k, R^C)) \rightarrow \text{Ext}_{C//B}^*(k, \text{Ext}_B^*(k, R))$$

is finite, so the spectral sequence

$$\text{Ext}_{C//B}^*(k, \text{Ext}_B^*(k, R)) \Rightarrow \text{Ext}_C^*(k, R)$$

is also a spectral sequence of finite modules over $R^C \otimes T$. □

The proof of A.11 is built out of a few special cases.

LEMMA A.12. *Suppose that E is a Hopf algebra of the form $E[x]$, where*

$$(A.13) \quad E[x] = \begin{cases} k[x]/x^2 & \text{if } |x| \text{ is odd} \\ k[x]/x^p & \text{if } |x| \text{ is even,} \end{cases}$$

and $R \rightarrow S$ is a finite map of Noetherian E -algebras. If the action of E on R is trivial, then

$$\text{Ext}_E^*(k, R) \rightarrow \text{Ext}_E^*(k, S)$$

is finite.

Proof. Let us take the case in which $E = k[x]/x^p$ with $|x|$ even. The others are similar. If M is an E -module, the cohomology

$$\text{Ext}_E^*(k, M)$$

is the cohomology of the complex

$$M \otimes \Lambda[a] \otimes k[b]$$

with differential

$$\begin{aligned} d(m \otimes b^k) &= x^{p-1}m \otimes a \otimes b^k, \\ d(m \otimes a \otimes b^k) &= xm \otimes b^{k+1}. \end{aligned}$$

When the action of E on M is trivial, the differential d is zero. The result now follows since the complex for calculating $\text{Ext}_E^*(k, S)$ is already a finite module over the $\text{Ext}_E^*(k, R)$. □

LEMMA A.14. *Suppose A is a Hopf algebra and $R \rightarrow S$ is a finite map of Noetherian A -algebras. If the action of A on R is trivial, then*

$$\text{Ext}_A^*(k, R) \rightarrow \text{Ext}_A^*(k, S)$$

is finite.

Proof. The proof is by induction on the dimension of A , the case in which the dimension of A is 1 being a tautology. Suppose that the dimension of A is greater than 1, and that the result is known to be true for Hopf algebras of dimension less than that of A .

Let $E \subseteq A$ be a central sub-Hopf algebra of the form (A.13), and let

$$\{E_r\}, \quad \text{and} \quad \{E'_r\}$$

be the associated Lyndon-Hochschild-Serre spectral sequences with coefficients in R and S , respectively. The spectral sequence $\{E_r\}$ is just the tensor product of R with the Lyndon-Hochschild-Serre spectral sequence with coefficients in k . By Theorem A.7 it is a spectral sequence of finite modules over a Noetherian ring of the form $R \otimes T$. It follows that the map

$$E_\infty \rightarrow E'_\infty$$

is finite, and so the map

$$\text{Ext}_A^*(k, R) \rightarrow \text{Ext}_A^*(k, S)$$

is finite by [4, Prop. 10.24]. □

Proof of Lemma A.9. Since S is finite over R and R is finite over R^A by Lemma A.8, S is finite over R^A . It follows from Lemma A.14 that

$$\text{Ext}_A^*(k, R^A) \rightarrow \text{Ext}_A^*(k, S)$$

is finite, so *a fortiori*

$$\text{Ext}_A^*(k, R) \rightarrow \text{Ext}_A^*(k, S)$$

is finite. This completes the proof. □

Proof of Theorem A.4. By Corollary A.11 we may assume that $R = k$. Choose an integer N with the property that x^{p^N} is invariant under the action of $C//B$ for every $x \in \text{Ext}_B(k, k)$, and let

$$S \subset \text{Ext}_B^*(k, k)$$

be the sub-algebra consisting of the $(p^N)^{\text{th}}$ powers of the elements in the image of $\text{Ext}_C^*(k, k)$. Then the maps

$$\begin{aligned} S &\rightarrow \text{Ext}_B^*(k, k), \quad \text{and} \\ \text{Ext}_{C//B}^*(k, S) &\rightarrow \text{Ext}_{C//B}^*(k, \text{Ext}_B^*(k, k)) \end{aligned}$$

are finite by A.8 and A.9. But

$$\text{Ext}_{C//B}^*(k, S) = \text{Ext}_{C//B}^*(k, k) \otimes S$$

is Noetherian, and consists of permanent cycles. Taking T to be this algebra establishes part iv), and completes the proof. \square

To deduce Theorem 4.13 requires

LEMMA A.15. *Suppose that $B \subseteq C$ is an inclusion of finite Hopf algebras. There is a sequence*

$$B = C_0 \triangleleft C_1 \cdots \triangleleft C_n = C$$

with each $C_i \triangleleft C_{i+1}$ normal, and with the property that the Hopf algebra C_{i+1}/C_i is of the form (A.13).

Proof. It suffices, by induction on $\dim_k C//B$, to show that if $B \neq C$ then there is a surjective map of Hopf algebras $C \rightarrow E[x]$ with the property that the composite

$$B \rightarrow C \rightarrow E[x]$$

is trivial (which means that it is the augmentation followed by the inclusion of the degree zero part). Since $B \neq C$ the map of dual algebras

$$C^* \rightarrow B^*$$

is not a monomorphism. It follows from [16, Prop. 3.9] that the map of primitives is not injective. Let D be a primitive in the kernel. The element D can be thought of as a derivation from A to k with the property that $D(b) = 0$ when $b \in B$.

Give x the degree $-|D|$. The map to $E[x]$ is then given by Taylor's formula:

$$a \mapsto \begin{cases} \sum_{n=0}^{p-1} D^n a \frac{x^n}{n!} & |D| \text{ even} \\ D^0 a + D(a) x & |D| \text{ odd.} \end{cases}$$

The powers of D are taken in the algebra C^* . In particular, D^0 , being the unit of C^* , is the augmentation. \square

Proof of Theorem 4.13. It suffices, by Lemma A.15 to deal with the case in which $B \subseteq C$ is normal. Let

$$b \in \text{Ext}_B^*(k, k)$$

be a cohomology class. By Lemma A.8 there is an integer M with the property that b^{p^M} is invariant under $C//B$. This gives a class in the E_2 -term of the Lyndon-Hochschild-Serre spectral sequence. For convenience, replace b^{p^M} with b . By Theorem A.4 there is an integer N with the property that $d_r b^{p^N} = 0$ for all r . The class in $\text{Ext}_C^*(k, k)$ represented by b^{p^N} is then the desired class. \square

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA

E-mail address: MJH@MATH.MIT.EDU

PURDUE UNIVERSITY, WEST LAFAYETTE, INDIANA

E-mail address: JHS@MATH.PURDUE.EDU

REFERENCES

- [1] J. F. ADAMS, On the structure and applications of the Steenrod algebra, *Comment. Math. Helv.* **32** (1958), 180–214.
- [2] ———, *Stable Homotopy and Generalised Homology*, University of Chicago Press, Chicago, 1974.
- [3] D. W. ANDERSON and D. M. DAVIS, A vanishing theorem in homological algebra, *Comment. Math. Helv.* **48** (1973), 318–327.
- [4] M. ATIYAH and I. MACDONALD, *Introduction to Commutative Algebra*, Addison-Wesley, Reading, Massachusetts, 1969.
- [5] N. BAAS, On bordism theory of manifolds with singularities, *Math. Scand.* **33** (1973), 279–302.
- [6] N. BAAS and I. MADSEN, On the realization of certain modules over the Steenrod algebra, *Math. Scand.* **31** (1972), 220–224.
- [7] M. G. BARRATT and M. E. MAHOWALD (eds.), Geometric applications of homotopy theory (Proc. Conf., Evanston, Ill., 1977), II, *Lecture Notes in Mathematics*, **658**, Berlin, Springer-Verlag, 1978.
- [8] N. BOURBAKI, *Commutative Algebra*, Elements of Mathematics, Addison-Wesley and Hermann, Paris, 1972.
- [9] A. K. BOUSFIELD, The Boolean algebra of spectra, *Comment. Math. Helv.* **54** (1979), 368–377.
- [10] A. K. BOUSFIELD and E. M. FRIEDLANDER, Homotopy theory of Γ -spaces, spectra, and bisimplicial sets, In: Barratt and Mahowald [7], pp. 80–130.
- [11] E. S. DEVINATZ, M. J. HOPKINS, and J. H. SMITH, Nilpotence and stable homotopy theory, *Ann. of Math.* **128** (1988), 207–241.
- [12] M. HAZEWINKEL, *Formal Groups and Applications*, Academic Press, New York, 1978.
- [13] M. J. HOPKINS, Global methods in homotopy theory, *Proc. 1985 LMS Symposium on Homotopy Theory* (J. D. S. Jones and E. Rees, eds.), 1987, pp. 73–96.
- [14] L. G. LEWIS, J. P. MAY, and M. STEINBERGER, *Equivariant Stable Homotopy Theory*, *Lecture Notes in Mathematics* **1213**, Springer-Verlag, New York, 1986.
- [15] H. R. MILLER and C. WILKERSON, Vanishing lines for modules over the Steenrod algebra, *J. Pure and App. Alg.* **22** (1981), 293–307.
- [16] J. W. MILNOR and J. C. MOORE, On the structure of Hopf algebras, *Ann. of Math.* **81** (1965), 211–264.
- [17] S. A. MITCHELL, Finite complexes with $A(n)$ -free cohomology, *Topology* **24** (1985), 227–246.
- [18] J. MORAVA, A product for the odd-primary bordism of manifolds with singularities, *Topology* **18** (1979), 177–186.
- [19] D. C. RAVENEL, Localization with respect to certain periodic homology theories, *Amer. J. Math.* **106** (1984), 351–414.
- [20] ———, *Complex Cobordism and Stable Homotopy Groups of Spheres*, Academic Press, Orlando, 1986.
- [21] ———, *Nilpotence and Periodicity in Stable Homotopy Theory*, *Ann. of Math. Studies*, vol. 128, Princeton University Press, Princeton, NJ, 1992.
- [22] C. A. ROBINSON, Obstruction theory and the strict associativity of Morava K -theories, *Advances in Homotopy* (Cambridge) (S. M. Salamon, B. Steer, and W. A. Sutherland, eds.), *London Mathematical Society Lecture Note Series*, vol. 139, Cambridge University Press, 1989, pp. 111–126.

- [23] W. M. SINGER, Steenrod squares in spectral sequences II, *Trans. A.M.S.* **175** (1973), 337–353.
- [24] J. SMITH, Finite complexes with vanishing lines of small slope, to appear.
- [25] C. WILKERSON, The cohomology algebras of finite dimensional Hopf algebras, *Trans. A.M.S.* **264** (1981), 137–150.

(Received July 9, 1992)