List of formulas for STK4011/9011 – Statistical Inference Theory (2025)

Chapter 2: Transformations and Expectations

Theorem 2.1.3 *Let X have cdf $F_X(x)$, let $Y = g(X)$, and let $\mathcal{X}$ and $\mathcal{Y}$ be defined as in (2.1.7).*

a. *If g is an increasing function on $\mathcal{X}$, $F_Y(y) = F_X\left(g^{-1}(y)\right)$ for $y \in \mathcal{Y}$.*

b. *If g is a decreasing function on $\mathcal{X}$ and X is a continuous random variable, $F_Y(y) = 1 - F_X\left(g^{-1}(y)\right)$ for $y \in \mathcal{Y}$.*

(2.1.7) $\mathcal{X} = \{x\colon\ f_X(x) > 0\}$ and $\mathcal{Y} = \{y\colon\ y = g(x) \text{ for some } x \in \mathcal{X}\}$.

Theorem 2.1.5 *Let X have pdf $f_X(x)$ and let $Y = g(X)$, where g is a monotone function. Let $\mathcal{X}$ and $\mathcal{Y}$ be defined by (2.1.7). Suppose that $f_X(x)$ is continuous on $\mathcal{X}$ and that $g^{-1}(y)$ has a continuous derivative on $\mathcal{Y}$. Then the pdf of Y is given by*

$$f_Y(y) = \begin{cases} f_X(g^{-1}(y))\left|\dfrac{d}{dy}g^{-1}(y)\right| & y \in \mathcal{Y} \\ 0 & \text{otherwise.} \end{cases} \tag{2.1.10}$$

Definition 2.2.1 The *expected value* or *mean* of a random variable $g(X)$, denoted by $\mathrm{E}\, g(X)$, is

$$\mathrm{E}\, g(X) = \begin{cases} \int_{-\infty}^{\infty} g(x) f_X(x)\, dx & \text{if } X \text{ is continuous} \\ \sum_{x \in \mathcal{X}} g(x) f_X(x) = \sum_{x \in \mathcal{X}} g(x) P(X = x) & \text{if } X \text{ is discrete,} \end{cases}$$

Theorem 2.2.5 *Let X be a random variable and let a, b, and c be constants. Then for any functions $g_1(x)$ and $g_2(x)$ whose expectations exist,*

a. $\mathrm{E}(ag_1(X) + bg_2(X) + c) = a\mathrm{E}\, g_1(X) + b\mathrm{E}\, g_2(X) + c$.

b. *If $g_1(x) \geq 0$ for all x, then $\mathrm{E}\, g_1(X) \geq 0$.*

c. *If $g_1(x) \geq g_2(x)$ for all x, then $\mathrm{E}\, g_1(X) \geq \mathrm{E}\, g_2(X)$.*

d. *If $a \leq g_1(x) \leq b$ for all x, then $a \leq \mathrm{E}\, g_1(X) \leq b$.*

Definition 2.3.1 For each integer n, the nth *moment of* X (or $F_X(x)$), μ_n', is

$$\mu_n' = \mathrm{E}\, X^n.$$

The nth *central moment of* X, μ_n, is

$$\mu_n = \mathrm{E}(X - \mu)^n,$$

where $\mu = \mu_1' = \mathrm{E}\, X$.

Definition 2.3.2 The *variance* of a random variable X is its second central moment, $\mathrm{Var}\, X = \mathrm{E}(X - \mathrm{E}\, X)^2$. The positive square root of $\mathrm{Var}\, X$ is the *standard deviation* of X.

Theorem 2.3.4 *If X is a random variable with finite variance, then for any constants a and b,*

$$\mathrm{Var}(aX + b) = a^2\, \mathrm{Var}\, X.$$

Definition 2.3.6 Let X be a random variable with cdf F_X. The *moment generating function (mgf) of* X (or F_X), denoted by $M_X(t)$, is

$$M_X(t) = \mathrm{E}\, e^{tX},$$

Theorem 2.3.7 *If X has mgf $M_X(t)$, then*

$$\mathrm{E}\, X^n = M_X^{(n)}(0),$$

where we define

$$M_X^{(n)}(0) = \frac{d^n}{dt^n} M_X(t)\bigg|_{t=0}.$$

That is, the nth moment is equal to the nth derivative of $M_X(t)$ evaluated at $t = 0$.

Theorem 2.3.11 *Let $F_X(x)$ and $F_Y(y)$ be two cdfs all of whose moments exist.*

a. *If X and Y have bounded support, then $F_X(u) = F_Y(u)$ for all u if and only if $\mathrm{E}\,X^r = \mathrm{E}\,Y^r$ for all integers $r = 0, 1, 2, \ldots$.*

b. *If the moment generating functions exist and $M_X(t) = M_Y(t)$ for all t in some neighborhood of 0, then $F_X(u) = F_Y(u)$ for all u.*

Theorem 2.3.12 (Convergence of mgfs) *Suppose $\{X_i,\ i = 1, 2, \ldots\}$ is a sequence of random variables, each with mgf $M_{X_i}(t)$. Furthermore, suppose that*

$$\lim_{i\to\infty} M_{X_i}(t) = M_X(t), \qquad \text{for all } t \text{ in a neighborhood of 0,}$$

and $M_X(t)$ is an mgf. Then there is a unique cdf F_X whose moments are determined by $M_X(t)$ and, for all x where $F_X(x)$ is continuous, we have

$$\lim_{i\to\infty} F_{X_i}(x) = F_X(x).$$

That is, convergence, for $|t| < h$, of mgfs to an mgf implies convergence of cdfs.

Theorem 2.3.15 *For any constants a and b, the mgf of the random variable $aX+b$ is given by*

$$M_{aX+b}(t) = e^{bt} M_X(at).$$

Chapter 3: Common Families of Distributions

A family of pdfs or pmfs is called an *exponential family* if it can be expressed as

$$f(x|\boldsymbol{\theta}) = h(x)c(\boldsymbol{\theta}) \exp\left(\sum_{i=1}^{k} w_i(\boldsymbol{\theta}) t_i(x)\right). \tag{3.4.1}$$

Here $h(x) \geq 0$ and $t_1(x), \ldots, t_k(x)$ are real-valued functions of the observation x (they cannot depend on $\boldsymbol{\theta}$), and $c(\boldsymbol{\theta}) \geq 0$ and $w_1(\boldsymbol{\theta}), \ldots, w_k(\boldsymbol{\theta})$ are real-valued functions of the possibly vector-valued parameter $\boldsymbol{\theta}$ (they cannot depend on x).

Definition 3.5.5 Let $f(x)$ be any pdf. Then for any μ, $-\infty < \mu < \infty$, and any $\sigma > 0$, the family of pdfs $(1/\sigma)f((x-\mu)/\sigma)$, indexed by the parameter (μ, σ), is called the *location–scale family with standard pdf $f(x)$*; μ is called the *location parameter* and σ is called the *scale parameter.*

Theorem 3.5.6 *Let $f(\cdot)$ be any pdf. Let μ be any real number, and let σ be any positive real number. Then X is a random variable with pdf $(1/\sigma)f((x-\mu)/\sigma)$ if and only if there exists a random variable Z with pdf $f(z)$ and $X = \sigma Z + \mu$.*

Theorem 3.6.1 (Chebychev's Inequality) *Let X be a random variable and let $g(x)$ be a nonnegative function. Then, for any $r > 0$,*

$$P(g(X) \geq r) \leq \frac{\mathrm{E}g(X)}{r}.$$

Chapter 4: Multiple Random Variables

If (X, Y) is a discrete bivariate random vector, then there is only a countable set of values for which the joint pmf of (X, Y) is positive. Call this set $\mathcal{A}$. Define the set $\mathcal{B} = \{(u, v) : u = g_1(x, y) \text{ and } v = g_2(x, y) \text{ for some } (x, y) \in \mathcal{A}\}$. Then $\mathcal{B}$ is the countable set of possible values for the discrete random vector (U, V). And if, for any $(u, v) \in \mathcal{B}$, A_{uv} is defined to be $\{(x, y) \in \mathcal{A} : g_1(x, y) = u \text{ and } g_2(x, y) = v\}$, then the joint pmf of (U, V), $f_{U,V}(u, v)$, can be computed from the joint pmf of (X, Y) by

$$f_{U,V}(u, v) = P(U = u, V = v) = P((X, Y) \in A_{uv}) = \sum_{(x,y)\in A_{uv}} f_{X,Y}(x, y). \tag{4.3.1}$$

If (X, Y) is a continuous random vector with joint pdf $f_{X,Y}(x, y)$, then the joint pdf of (U, V) can be expressed in terms of $f_{X,Y}(x, y)$ in a manner analogous to (2.1.8). As before, $\mathcal{A} = \{(x, y) : f_{X,Y}(x, y) > 0\}$ and $\mathcal{B} = \{(u, v) : u = g_1(x, y) \text{ and } v = g_2(x, y) \text{ for some } (x, y) \in \mathcal{A}\}$. The joint pdf $f_{U,V}(u, v)$ will be positive on the set $\mathcal{B}$. For the simplest version of this result we assume that the transformation $u = g_1(x, y)$ and $v = g_2(x, y)$ defines a one-to-one transformation of $\mathcal{A}$ onto $\mathcal{B}$. The transformation is onto because of the definition of $\mathcal{B}$. We are assuming that for each $(u, v) \in \mathcal{B}$ there is only one $(x, y) \in \mathcal{A}$ such that $(u, v) = (g_1(x, y), g_2(x, y))$. For such a one-to-one, onto transformation, we can solve the equations $u = g_1(x, y)$ and $v = g_2(x, y)$ for x and y in terms of u and v. We will denote this inverse transformation by $x = h_1(u, v)$ and $y = h_2(u, v)$. The role played by a derivative in the univariate case is now played by a quantity called the *Jacobian of the transformation.* This function of (u, v), denoted by J, is the *determinant of a matrix* of partial derivatives. It is defined by

$$J = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial x}{\partial u}\frac{\partial y}{\partial v} - \frac{\partial y}{\partial u}\frac{\partial x}{\partial v},$$

where

$$\frac{\partial x}{\partial u} = \frac{\partial h_1(u,v)}{\partial u}, \quad \frac{\partial x}{\partial v} = \frac{\partial h_1(u,v)}{\partial v}, \quad \frac{\partial y}{\partial u} = \frac{\partial h_2(u,v)}{\partial u}, \quad \text{and} \quad \frac{\partial y}{\partial v} = \frac{\partial h_2(u,v)}{\partial v}.$$

We assume that J is not identically 0 on $\mathcal{B}$. Then the joint pdf of (U,V) is 0 outside the set $\mathcal{B}$ and on the set $\mathcal{B}$ is given by

$$f_{U,V}(u,v) = f_{X,Y}(h_1(u,v),\ h_2(u,v))|J|, \tag{4.3.2}$$

Lemma 4.2.7 *Let (X,Y) be a bivariate random vector with joint pdf or pmf $f(x,y)$. Then X and Y are independent random variables if and only if there exist functions $g(x)$ and $h(y)$ such that, for every $x \in \Re$ and $y \in \Re$,*

$$f(x,y) = g(x)h(y).$$

Theorem 4.2.12 *Let X and Y be independent random variables with moment generating functions $M_X(t)$ and $M_Y(t)$. Then the moment generating function of the random variable $Z = X + Y$ is given by*

$$M_Z(t) = M_X(t)M_Y(t).$$

Theorem 4.3.5 *Let X and Y be independent random variables. Let $g(x)$ be a function only of x and $h(y)$ be a function only of y. Then the random variables $U = g(X)$ and $V = h(Y)$ are independent.*

Theorem 4.4.3 *If X and Y are any two random variables, then*

$$\mathrm{E}X = \mathrm{E}\left(\mathrm{E}(X|Y)\right), \tag{4.4.1}$$

provided that the expectations exist.

Theorem 4.4.7 (Conditional variance identity) *For any two random variables X and Y,*

$$\mathrm{Var}\, X = \mathrm{E}\left(\mathrm{Var}(X|Y)\right) + \mathrm{Var}\left(\mathrm{E}(X|Y)\right), \tag{4.4.4}$$

provided that the expectations exist.

Definition 4.5.1 The *covariance of X and Y* is the number defined by

$$\mathrm{Cov}(X,Y) = \mathrm{E}\left((X-\mu_X)(Y-\mu_Y)\right).$$

Theorem 4.5.5 *If X and Y are independent random variables, then* $\mathrm{Cov}(X,Y) = 0$ *and* $\rho_{XY} = 0$.

Theorem 4.5.6 *If X and Y are any two random variables and a and b are any two constants, then*

$$\mathrm{Var}(aX+bY) = a^2\mathrm{Var}\, X + b^2\mathrm{Var}\, Y + 2ab\mathrm{Cov}(X,Y).$$

If X and Y are independent random variables, then

$$\mathrm{Var}(aX+bY) = a^2\mathrm{Var}\, X + b^2\mathrm{Var}\, Y.$$

Theorem 4.6.7 (Generalization of Theorem 4.2.12) *Let $X_1,\ldots,X_n$ be mutually independent random variables with mgfs $M_{X_1}(t),\ldots,M_{X_n}(t)$. Let $Z = X_1+\cdots+X_n$. Then the mgf of Z is*

$$M_Z(t) = M_{X_1}(t)\cdot\cdots\cdot M_{X_n}(t).$$

In particular, if $X_1,\ldots,X_n$ all have the same distribution with mgf $M_X(t)$, then

$$M_Z(t) = \left(M_X(t)\right)^n.$$

Theorem 4.7.7 (Jensen's Inequality) *For any random variable X, if $g(x)$ is a convex function, then*

$$\mathrm{E}g(X) \geq g(\mathrm{E}X).$$

Equality holds if and only if, for every line $a+bx$ that is tangent to $g(x)$ at $x = EX$, $P(g(X) = a+bX) = 1$.

Chapter 5: Multiple Random Variables

Theorem 5.2.6 *Let $X_1,\ldots,X_n$ be a random sample from a population with mean μ and variance $\sigma^2 < \infty$. Then*

a. $\mathrm{E}\bar{X} = \mu$,

b. $\mathrm{Var}\,\bar{X} = \dfrac{\sigma^2}{n}$,

c. $\mathrm{E}S^2 = \sigma^2$.

Theorem 5.2.7 *Let $X_1, \ldots, X_n$ be a random sample from a population with mgf $M_X(t)$. Then the mgf of the sample mean is*

$$M_{\bar{X}}(t) = [M_X(t/n)]^n.$$

Theorem 5.2.9 *If X and Y are independent continuous random variables with pdfs $f_X(x)$ and $f_Y(y)$, then the pdf of $Z = X + Y$ is*

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(w) f_Y(z - w)\, dw. \tag{5.2.3}$$

Theorem 5.4.3 *Let $X_1, \ldots, X_n$ be a random sample from a discrete distribution with pmf $f_X(x_i) = p_i$, where $x_1 < x_2 < \cdots$ are the possible values of X in ascending*

Theorem 5.4.4 *Let $X_{(1)}, \ldots, X_{(n)}$ denote the order statistics of a random sample, $X_1, \ldots, X_n$, from a continuous population with cdf $F_X(x)$ and pdf $f_X(x)$. Then the pdf of $X_{(j)}$ is*

$$f_{X_{(j)}}(x) = \frac{n!}{(j-1)!(n-j)!} f_X(x)[F_X(x)]^{j-1}[1 - F_X(x)]^{n-j}. \tag{5.4.4}$$

Theorem 5.4.6 *Let $X_{(1)}, \ldots, X_{(n)}$ denote the order statistics of a random sample, $X_1, \ldots, X_n$, from a continuous population with cdf $F_X(x)$ and pdf $f_X(x)$. Then the joint pdf of $X_{(i)}$ and $X_{(j)}, 1 \le i < j \le n$, is*

$$f_{X_{(i)},X_{(j)}}(u, v) = \frac{n!}{(i-1)!(j-1-i)!(n-j)!} f_X(u) f_X(v)[F_X(u)]^{i-1} \times [F_X(v) - F_X(u)]^{j-1-i}[1 - F_X(v)]^{n-j} \tag{5.4.7}$$

for $-\infty < u < v < \infty$.

Definition 5.5.1 A sequence of random variables, $X_1, X_2, \ldots$, *converges in probability* to a random variable X if, for every $\epsilon > 0$,

$$\lim_{n\to\infty} P(|X_n - X| \ge \epsilon) = 0 \quad \text{or, equivalently,} \quad \lim_{n\to\infty} P(|X_n - X| < \epsilon) = 1.$$

Theorem 5.5.2 (Weak Law of Large Numbers) *Let $X_1, X_2, \ldots$ be iid random variables with $\mathrm{E}X_i = \mu$ and $\mathrm{Var}\, X_i = \sigma^2 < \infty$. Define $\bar{X}_n = (1/n)\sum_{i=1}^n X_i$. Then, for every $\epsilon > 0$,*

$$\lim_{n\to\infty} P(|\bar{X}_n - \mu| < \epsilon) = 1;$$

that is, $\bar{X}_n$ converges in probability to μ.

Theorem 5.5.4 *Suppose that $X_1, X_2, \ldots$ converges in probability to a random variable X and that h is a continuous function. Then $h(X_1), h(X_2), \ldots$ converges in probability to $h(X)$.*

Definition 5.5.6 A sequence of random variables, $X_1, X_2, \ldots$, *converges almost surely* to a random variable X if, for every $\epsilon > 0$,

$$P(\lim_{n\to\infty} |X_n - X| < \epsilon) = 1.$$

Theorem 5.5.9 (Strong Law of Large Numbers) *Let $X_1, X_2, \ldots$ be iid random variables with $\mathrm{E}X_i = \mu$ and $\mathrm{Var}\, X_i = \sigma^2 < \infty$, and define $\bar{X}_n = (1/n)\sum_{i=1}^n X_i$. Then, for every $\epsilon > 0$,*

$$P(\lim_{n\to\infty} |\bar{X}_n - \mu| < \epsilon) = 1;$$

that is, $\bar{X}_n$ converges almost surely to μ.

Definition 5.5.10 A sequence of random variables, $X_1, X_2, \ldots$, *converges in distribution* to a random variable X if

$$\lim_{n\to\infty} F_{X_n}(x) = F_X(x)$$

at all points x where $F_X(x)$ is continuous.

Theorem 5.5.12 *If the sequence of random variables, $X_1, X_2, \ldots$, converges in probability to a random variable X, the sequence also converges in distribution to X.*

Theorem 5.5.13 *The sequence of random variables, $X_1, X_2, \ldots$, converges in probability to a constant μ if and only if the sequence also converges in distribution to μ. That is, the statement*

$$P(|X_n - \mu| > \varepsilon) \to 0 \text{ for every } \varepsilon > 0$$

is equivalent to

$$P(X_n \le x) \to \begin{cases} 0 & \text{if } x < \mu \\ 1 & \text{if } x > \mu. \end{cases}$$

Theorem 5.5.15 (Stronger form of the Central Limit Theorem) *Let $X_1, X_2, \ldots$ be a sequence of iid random variables with $\mathrm{E}X_i = \mu$ and $0 < \mathrm{Var}\, X_i = \sigma^2 < \infty$. Define $\bar{X}_n = (1/n)\sum_{i=1}^n X_i$. Let $G_n(x)$ denote the cdf of $\sqrt{n}(\bar{X}_n - \mu)/\sigma$. Then, for any x, $-\infty < x < \infty$,*

$$\lim_{n\to\infty} G_n(x) = \int_{-\infty}^{x} \frac{1}{\sqrt{2\pi}} e^{-y^2/2}\, dy;$$

that is, $\sqrt{n}(\bar{X}_n - \mu)/\sigma$ has a limiting standard normal distribution.

Theorem 5.5.17 (Slutsky's Theorem) *If $X_n \to X$ in distribution and $Y_n \to a$, a constant, in probability, then*

a. *$Y_n X_n \to aX$ in distribution.*

b. *$X_n + Y_n \to X + a$ in distribution.*

Theorem 5.5.24 (Delta Method) *Let Y_n be a sequence of random variables that satisfies $\sqrt{n}(Y_n - \theta) \to \mathrm{n}(0, \sigma^2)$ in distribution. For a given function g and a specific value of θ, suppose that $g'(\theta)$ exists and is not 0. Then*

(5.5.10) $$\sqrt{n}[g(Y_n) - g(\theta)] \to \mathrm{n}(0, \sigma^2[g'(\theta)]^2) \text{ in distribution.}$$

Theorem 5.5.26 (Second-order Delta Method) *Let Y_n be a sequence of random variables that satisfies $\sqrt{n}(Y_n - \theta) \to \mathrm{n}(0, \sigma^2)$ in distribution. For a given function g and a specific value of θ, suppose that $g'(\theta) = 0$ and $g''(\theta)$ exists and is not 0. Then*

(5.5.13) $$n[g(Y_n) - g(\theta)] \to \sigma^2 \frac{g''(\theta)}{2} \chi_1^2 \text{ in distribution.}$$

Theorem 5.5.28 (Multivariate Delta Method) *Let $\mathbf{X}_1, \ldots, \mathbf{X}_n$ be a random sample with $\mathrm{E}(X_{ij}) = \mu_i$ and $\mathrm{Cov}(X_{ik}, X_{jk}) = \sigma_{ij}$. For a given function g with continuous first partial derivatives and a specific value of $\boldsymbol{\mu} = (\mu_1, \ldots, \mu_p)$ for which $\tau^2 = \Sigma\Sigma\sigma_{ij} \frac{\partial g(\boldsymbol{\mu})}{\partial \mu_i} \cdot \frac{\partial g(\boldsymbol{\mu})}{\partial \mu_j} > 0$,*

$$\sqrt{n}[g(\bar{X}_1, \ldots, \bar{X}_s) - g(\mu_1, \ldots, \mu_p)] \to \mathrm{n}(0, \tau^2) \text{ in distribution .}$$

Chapter 6: Principles of Data Reduction

Definition 6.2.1 A statistic $T(\mathbf{X})$ is a *sufficient statistic for θ* if the conditional distribution of the sample $\mathbf{X}$ given the value of $T(\mathbf{X})$ does not depend on θ.

Theorem 6.2.2 *If $p(\mathbf{x}|\theta)$ is the joint pdf or pmf of $\mathbf{X}$ and $q(t|\theta)$ is the pdf or pmf of $T(\mathbf{X})$, then $T(\mathbf{X})$ is a sufficient statistic for θ if, for every $\mathbf{x}$ in the sample space, the ratio $p(\mathbf{x}|\theta)/q(T(\mathbf{x})|\theta)$ is constant as a function of θ.*

Theorem 6.2.6 (Factorization Theorem) *Let $f(\mathbf{x}|\theta)$ denote the joint pdf or pmf of a sample $\mathbf{X}$. A statistic $T(\mathbf{X})$ is a sufficient statistic for θ if and only if there exist functions $g(t|\theta)$ and $h(\mathbf{x})$ such that, for all sample points $\mathbf{x}$ and all parameter points θ,*

(6.2.3) $$f(\mathbf{x}|\theta) = g(T(\mathbf{x})|\theta)h(\mathbf{x}).$$

Theorem 6.2.10 *Let $X_1, \ldots, X_n$ be iid observations from a pdf or pmf $f(x|\boldsymbol{\theta})$ that belongs to an exponential family given by*

$$f(x|\boldsymbol{\theta}) = h(x)c(\boldsymbol{\theta}) \exp\left(\sum_{i=1}^{k} w_i(\boldsymbol{\theta}) t_i(x)\right),$$

where $\boldsymbol{\theta} = (\theta_1, \theta_2, \ldots, \theta_d)$, $d \le k$. Then

$$T(\mathbf{X}) = \left(\sum_{j=1}^{n} t_1(X_j), \ldots, \sum_{j=1}^{n} t_k(X_j)\right)$$

is a sufficient statistic for $\boldsymbol{\theta}$.

Definition 6.2.11 A sufficient statistic $T(\mathbf{X})$ is called a *minimal sufficient statistic* if, for any other sufficient statistic $T'(\mathbf{X})$, $T(\mathbf{x})$ is a function of $T'(\mathbf{x})$.

Theorem 6.2.13 *Let $f(\mathbf{x}|\theta)$ be the pmf or pdf of a sample $\mathbf{X}$. Suppose there exists a function $T(\mathbf{x})$ such that, for every two sample points $\mathbf{x}$ and $\mathbf{y}$, the ratio $f(\mathbf{x}|\theta)/f(\mathbf{y}|\theta)$ is constant as a function of θ if and only if $T(\mathbf{x}) = T(\mathbf{y})$. Then $T(\mathbf{X})$ is a minimal sufficient statistic for θ.*

Definition 6.2.21 Let $f(t|\theta)$ be a family of pdfs or pmfs for a statistic $T(\mathbf{X})$. The family of probability distributions is called *complete* if $\mathrm{E}_\theta g(T) = 0$ for all θ implies $P_\theta(g(T) = 0) = 1$ for all θ. Equivalently, $T(\mathbf{X})$ is called a *complete statistic.*

Theorem 6.2.24 (Basu's Theorem) *If $T(\mathbf{X})$ is a complete and minimal sufficient statistic, then $T(\mathbf{X})$ is independent of every ancillary statistic.*

Theorem 6.2.25 (Complete statistics in the exponential family) *Let $X_1, \ldots, X_n$ be iid observations from an exponential family with pdf or pmf of the form*

$$f(x|\boldsymbol{\theta}) = h(x)c(\boldsymbol{\theta}) \exp\left(\sum_{j=1}^{k} w(\theta_j)t_j(x)\right), \tag{6.2.7}$$

where $\boldsymbol{\theta} = (\theta_1, \theta_2, \ldots, \theta_k)$. Then the statistic

$$T(\mathbf{X}) = \left(\sum_{i=1}^{n} t_1(X_i), \sum_{i=1}^{n} t_2(X_i), \ldots, \sum_{i=1}^{n} t_k(X_i)\right)$$

is complete as long as the parameter space Θ contains an open set in $\Re^k$.

Theorem 6.2.28 *If a minimal sufficient statistic exists, then any complete statistic is also a minimal sufficient statistic.*

Chapter 7: Point Estimation

Theorem 7.2.10 (Invariance property of MLEs) *If $\hat{\theta}$ is the MLE of θ, then for any function $\tau(\theta)$, the MLE of $\tau(\theta)$ is $\tau(\hat{\theta})$.*

Definition 7.3.7 An estimator W^* is a *best unbiased estimator* of $\tau(\theta)$ if it satisfies $\mathrm{E}_\theta W^* = \tau(\theta)$ for all θ and, for any other estimator W with $\mathrm{E}_\theta W = \tau(\theta)$, we have $\mathrm{Var}_\theta W^* \leq \mathrm{Var}_\theta W$ for all θ. W^* is also called a *uniform minimum variance unbiased estimator* (UMVUE) of $\tau(\theta)$.

Theorem 7.3.9 (Cramér–Rao Inequality) *Let $X_1, \ldots, X_n$ be a sample with pdf $f(\mathbf{x}|\theta)$, and let $W(\mathbf{X}) = W(X_1, \ldots, X_n)$ be any estimator satisfying*

$$\frac{d}{d\theta}\mathrm{E}_\theta W(\mathbf{X}) = \int_{\mathcal{X}} \frac{\partial}{\partial\theta}\left[W(\mathbf{x})f(\mathbf{x}|\theta)\right] d\mathbf{x}$$

(7.3.4) *and*

$$\mathrm{Var}_\theta W(\mathbf{X}) < \infty.$$

Then

$$\mathrm{Var}_\theta\left(W(\mathbf{X})\right) \geq \frac{\left(\frac{d}{d\theta}\mathrm{E}_\theta W(\mathbf{X})\right)^2}{\mathrm{E}_\theta\left(\left(\frac{\partial}{\partial\theta}\log f(\mathbf{X}|\theta)\right)^2\right)}. \tag{7.3.5}$$

Corollary 7.3.10 (Cramér–Rao Inequality, iid case) *If the assumptions of Theorem 7.3.9 are satisfied and, additionally, if $X_1, \ldots, X_n$ are iid with pdf $f(x|\theta)$, then*

$$\mathrm{Var}_\theta W(\mathbf{X}) \geq \frac{\left(\frac{d}{d\theta}\mathrm{E}_\theta W(\mathbf{X})\right)^2}{n\mathrm{E}_\theta\left(\left(\frac{\partial}{\partial\theta}\log f(X|\theta)\right)^2\right)}.$$

Lemma 7.3.11 *If $f(x|\theta)$ satisfies*

$$\frac{d}{d\theta}\mathrm{E}_\theta\left(\frac{\partial}{\partial\theta}\log f(X|\theta)\right) = \int \frac{\partial}{\partial\theta}\left[\left(\frac{\partial}{\partial\theta}\log f(x|\theta)\right)f(x|\theta)\right] dx$$

(true for an exponential family), then

$$\mathrm{E}_\theta\left(\left(\frac{\partial}{\partial\theta}\log f(X|\theta)\right)^2\right) = -\mathrm{E}_\theta\left(\frac{\partial^2}{\partial\theta^2}\log f(X|\theta)\right).$$

Corollary 7.3.15 (Attainment) *Let $X_1, \ldots, X_n$ be iid $f(x|\theta)$, where $f(x|\theta)$ satisfies the conditions of the Cramér–Rao Theorem. Let $L(\theta|\mathbf{x}) = \prod_{i=1}^n f(x_i|\theta)$ denote the likelihood function. If $W(\mathbf{X}) = W(X_1, \ldots, X_n)$ is any unbiased estimator of $\tau(\theta)$, then $W(\mathbf{X})$ attains the Cramér–Rao Lower Bound if and only if*

$$a(\theta)[W(\mathbf{x}) - \tau(\theta)] = \frac{\partial}{\partial\theta}\log L(\theta|\mathbf{x}) \tag{7.3.12}$$

for some function $a(\theta)$.

Theorem 7.3.17 (Rao–Blackwell) *Let W be any unbiased estimator of $\tau(\theta)$, and let T be a sufficient statistic for θ. Define $\phi(T) = \mathrm{E}(W|T)$. Then $\mathrm{E}_\theta\phi(T) = \tau(\theta)$ and $\mathrm{Var}_\theta\,\phi(T) \leq \mathrm{Var}_\theta W$ for all θ; that is, $\phi(T)$ is a uniformly better unbiased estimator of $\tau(\theta)$.*

Theorem 7.3.19 *If W is a best unbiased estimator of $\tau(\theta)$, then W is unique.*

Theorem 7.3.20 *If $\mathrm{E}_\theta W = \tau(\theta), W$ is the best unbiased estimator of $\tau(\theta)$ if and only if W is uncorrelated with all unbiased estimators of 0.*

Theorem 7.3.23 *Let T be a complete sufficient statistic for a parameter θ, and let $\phi(T)$ be any estimator based only on T. Then $\phi(T)$ is the unique best unbiased estimator of its expected value.*

Chapter 8: Hypothesis Testing

Definition 8.2.1 The *likelihood ratio test statistic* for testing $H_0: \theta \in \Theta_0$ versus $H_1: \theta \in \Theta_0^c$ is

$$\lambda(\mathbf{x}) = \frac{\sup_{\Theta_0} L(\theta|\mathbf{x})}{\sup_{\Theta} L(\theta|\mathbf{x})}.$$

A *likelihood ratio test* (LRT) is any test that has a rejection region of the form $\{\mathbf{x}: \lambda(\mathbf{x}) \leq c\}$, where c is any number satisfying $0 \leq c \leq 1$.

Theorem 8.2.4 *If $T(\mathbf{X})$ is a sufficient statistic for θ and $\lambda^*(t)$ and $\lambda(\mathbf{x})$ are the LRT statistics based on T and $\mathbf{X}$, respectively, then $\lambda^*(T(\mathbf{x})) = \lambda(\mathbf{x})$ for every $\mathbf{x}$ in the sample space.*

Definition 8.3.1 The *power function* of a hypothesis test with rejection region R is the function of θ defined by $\beta(\theta) = P_\theta(\mathbf{X} \in R)$.

Definition 8.3.5 For $0 \leq \alpha \leq 1$, a test with power function $\beta(\theta)$ is a *size α test* if $\sup_{\theta \in \Theta_0} \beta(\theta) = \alpha$.

Definition 8.3.6 For $0 \leq \alpha \leq 1$, a test with power function $\beta(\theta)$ is a *level α test* if $\sup_{\theta \in \Theta_0} \beta(\theta) \leq \alpha$.

Definition 8.3.9 A test with power function $\beta(\theta)$ is *unbiased* if $\beta(\theta') \geq \beta(\theta'')$ for every $\theta' \in \Theta_0^c$ and $\theta'' \in \Theta_0$.

Definition 8.3.11 Let $\mathcal{C}$ be a class of tests for testing $H_0: \theta \in \Theta_0$ versus $H_1: \theta \in \Theta_0^c$. A test in class $\mathcal{C}$, with power function $\beta(\theta)$, is a *uniformly most powerful* (UMP) *class $\mathcal{C}$ test* if $\beta(\theta) \geq \beta'(\theta)$ for every $\theta \in \Theta_0^c$ and every $\beta'(\theta)$ that is a power function of a test in class $\mathcal{C}$.

Theorem 8.3.12 (Neyman–Pearson Lemma) *Consider testing $H_0: \theta = \theta_0$ versus $H_1: \theta = \theta_1$, where the pdf or pmf corresponding to θ_i is $f(\mathbf{x}|\theta_i), i = 0, 1$, using a test with rejection region R that satisfies*

$$\mathbf{x} \in R \text{ if } f(\mathbf{x}|\theta_1) > k f(\mathbf{x}|\theta_0)$$

(8.3.1) *and*

$$\mathbf{x} \in R^c \text{ if } f(\mathbf{x}|\theta_1) < k f(\mathbf{x}|\theta_0),$$

for some $k \geq 0$, and

$$\alpha = P_{\theta_0}(\mathbf{X} \in R). \quad (8.3.2)$$

Then

a. *(Sufficiency) Any test that satisfies (8.3.1) and (8.3.2) is a UMP level α test.*

b. *(Necessity) If there exists a test satisfying (8.3.1) and (8.3.2) with $k > 0$, then every UMP level α test is a size α test (satisfies (8.3.2)) and every UMP level α test satisfies (8.3.1) except perhaps on a set A satisfying $P_{\theta_0}(\mathbf{X} \in A) = P_{\theta_1}(\mathbf{X} \in A) = 0$.*

Corollary 8.3.13 *Consider the hypothesis problem posed in Theorem 8.3.12. Suppose $T(\mathbf{X})$ is a sufficient statistic for θ and $g(t|\theta_i)$ is the pdf or pmf of T corresponding to θ_i, $i = 0, 1$. Then any test based on T with rejection region S (a subset of the sample space of T) is a UMP level α test if it satisfies*

$$t \in S \text{ if } g(t|\theta_1) > k g(t|\theta_0)$$

(8.3.4) *and*

$$t \in S^c \text{ if } g(t|\theta_1) < k g(t|\theta_0),$$

for some $k \geq 0$, where

$$\alpha = P_{\theta_0}(T \in S). \quad (8.3.5)$$

Definition 8.3.16 A family of pdfs or pmfs $\{g(t|\theta): \theta \in \Theta\}$ for a univariate random variable T with real-valued parameter θ has a *monotone likelihood ratio* (MLR) if, for every $\theta_2 > \theta_1$, $g(t|\theta_2)/g(t|\theta_1)$ is a monotone (nonincreasing or nondecreasing) function of t on $\{t: g(t|\theta_1) > 0 \text{ or } g(t|\theta_2) > 0\}$. Note that $c/0$ is defined as ∞ if $0 < c$.

Theorem 8.3.17 (Karlin–Rubin) *Consider testing $H_0: \theta \leq \theta_0$ versus $H_1: \theta > \theta_0$. Suppose that T is a sufficient statistic for θ and the family of pdfs or pmfs $\{g(t|\theta): \theta \in \Theta\}$ of T has an MLR.* Then for any t_0, the test that rejects H_0 if and only if $T > t_0$ is a UMP level α test, where $\alpha = P_{\theta_0}(T > t_0)$.* *Assumes nondecreasing LR.

Chapter 9: Interval Estimation

Definition 9.1.1 An *interval estimate* of a real-valued parameter θ is any pair of functions, $L(x_1, \ldots, x_n)$ and $U(x_1, \ldots, x_n)$, of a sample that satisfy $L(\mathbf{x}) \leq U(\mathbf{x})$ for all $\mathbf{x} \in \mathcal{X}$. If $\mathbf{X} = \mathbf{x}$ is observed, the inference $L(\mathbf{x}) \leq \theta \leq U(\mathbf{x})$ is made. The random interval $[L(\mathbf{X}), U(\mathbf{X})]$ is called an *interval estimator*.

Definition 9.1.4 For an interval estimator $[L(\mathbf{X}), U(\mathbf{X})]$ of a parameter θ, the *coverage probability* of $[L(\mathbf{X}), U(\mathbf{X})]$ is the probability that the random interval $[L(\mathbf{X}), U(\mathbf{X})]$ covers the true parameter, θ. In symbols, it is denoted by either $P_\theta(\theta \in [L(\mathbf{X}), U(\mathbf{X})])$ or $P(\theta \in [L(\mathbf{X}), U(\mathbf{X})]|\theta)$.

Definition 9.1.5 For an interval estimator $[L(\mathbf{X}), U(\mathbf{X})]$ of a parameter θ, the *confidence coefficient* of $[L(\mathbf{X}), U(\mathbf{X})]$ is the infimum of the coverage probabilities, $\inf_\theta P_\theta(\theta \in [L(\mathbf{X}), U(\mathbf{X})])$.

Theorem 9.2.2 *For each $\theta_0 \in \Theta$, let $A(\theta_0)$ be the acceptance region of a level α test of $H_0: \theta = \theta_0$. For each $\mathbf{x} \in \mathcal{X}$, define a set $C(\mathbf{x})$ in the parameter space by*

$$C(\mathbf{x}) = \{\theta_0 : \mathbf{x} \in A(\theta_0)\}. \tag{9.2.1}$$

Then the random set $C(\mathbf{X})$ is a $1-\alpha$ confidence set. Conversely, let $C(\mathbf{X})$ be a $1-\alpha$ confidence set. For any $\theta_0 \in \Theta$, define

$$A(\theta_0) = \{\mathbf{x} : \theta_0 \in C(\mathbf{x})\}.$$

Then $A(\theta_0)$ is the acceptance region of a level α test of $H_0: \theta = \theta_0$.

Chapter 10: Asymptotic Evaluations

Definition 10.1.1 A sequence of estimators $W_n = W_n(X_1, \ldots, X_n)$ is a *consistent sequence of estimators* of the parameter θ if, for every $\epsilon > 0$ and every $\theta \in \Theta$,

$$\lim_{n\to\infty} P_\theta(|W_n - \theta| < \epsilon) = 1. \tag{10.1.1}$$

Theorem 10.1.3 *If W_n is a sequence of estimators of a parameter θ satisfying*

i. $\lim_{n\to\infty} \mathrm{Var}_\theta W_n = 0$,

ii. $\lim_{n\to\infty} \mathrm{Bias}_\theta W_n = 0$,

for every $\theta \in \Theta$, then W_n is a consistent sequence of estimators of θ.

Definition 10.1.9 For an estimator T_n, suppose that $k_n(T_n - \tau(\theta)) \to \mathrm{n}(0, \sigma^2)$ in distribution. The parameter σ^2 is called the *asymptotic variance* or *variance of the limit distribution* of T_n.

Definition 10.1.11 A sequence of estimators W_n is *asymptotically efficient* for a parameter $\tau(\theta)$ if $\sqrt{n}[W_n - \tau(\theta)] \to \mathrm{n}[0, v(\theta)]$ in distribution and

$$v(\theta) = \frac{[\tau'(\theta)]^2}{\mathrm{E}_\theta\left(\left(\frac{\partial}{\partial\theta}\log f(X|\theta)\right)^2\right)};$$

that is, the asymptotic variance of W_n achieves the Cramér–Rao Lower Bound.

Theorem 10.1.12 (Asymptotic efficiency of MLEs) *Let $X_1, X_2, \ldots,$ be iid $f(x|\theta)$, let $\hat{\theta}$ denote the MLE of θ, and let $\tau(\theta)$ be a continuous function of θ. Under the regularity conditions in Miscellanea 10.6.2 on $f(x|\theta)$ and, hence, $L(\theta|\mathbf{x})$,*

$$\sqrt{n}[\tau(\hat{\theta}) - \tau(\theta)] \to \mathrm{n}[0, v(\theta)],$$

where $v(\theta)$ is the Cramér–Rao Lower Bound. That is, $\tau(\hat{\theta})$ is a consistent and asymptotically efficient estimator of $\tau(\theta)$.

Definition 10.1.16 If two estimators W_n and V_n satisfy

$$\sqrt{n}[W_n - \tau(\theta)] \to \mathrm{n}[0, \sigma_W^2]$$
$$\sqrt{n}[V_n - \tau(\theta)] \to \mathrm{n}[0, \sigma_V^2]$$

in distribution, the *asymptotic relative efficiency* (ARE) of V_n with respect to W_n is

$$\mathrm{ARE}(V_n, W_n) = \frac{\sigma_W^2}{\sigma_V^2}.$$

Theorem 10.3.1 (Asymptotic distribution of the LRT—simple H_0) *For testing $H_0 : \theta = \theta_0$ versus $H_1 : \theta \neq \theta_0$, suppose $X_1, \ldots, X_n$ are iid $f(x|\theta)$, $\hat{\theta}$ is the MLE of θ, and $f(x|\theta)$ satisfies the regularity conditions in Miscellanea 10.6.2. Then under H_0, as $n \to \infty$,*

$$-2\log\lambda(\mathbf{X}) \to \chi_1^2 \text{ in distribution,}$$

where χ_1^2 is a χ^2 random variable with 1 degree of freedom.

Theorem 10.3.3 *Let $X_1, \ldots, X_n$ be a random sample from a pdf or pmf $f(x|\theta)$. Under the regularity conditions in Miscellanea 10.6.2, if $\theta \in \Theta_0$, then the distribution of the statistic $-2\log\lambda(\mathbf{X})$ converges to a chi squared distribution as the sample size $n \to \infty$. The degrees of freedom of the limiting distribution is the difference between the number of free parameters specified by $\theta \in \Theta_0$ and the number of free parameters specified by $\theta \in \Theta$.*