

ECON 4130  
HG revised Oct 10

## Lecture Notes to Rice Chapter 8

*On how to handle inference in models with more than one unknown parameter when the limit distribution for the estimators is multivariate normal.*

### 0 Introduction - summary of one-parameter (asymptotic) inference based on estimators that are asymptotically normally distributed.

Let  $\theta$  be a parameter of interest, the true value of which being unknown, in an econometric model. We wish to perform inference on  $\theta$  based on a data set. Let  $\hat{\theta} = \hat{\theta}_n$  be an estimator of  $\theta$  based on  $n$  observations or  $n$  observation vectors. Inference on  $\theta$  based on the exact distribution of  $\hat{\theta}$  is only possible in some situations. More often than not we need to resort to approximate methods - e.g., asymptotic methods for large or moderately large samples or simulation techniques (e.g. bootstrap) in small or moderately large samples.

Very often (e.g. in most cases of mle, mme, ols, gls, etc estimators for cross-section data and quite often for panel- and time series data as well) we have from theory some theorem that says

$$(0-1) \quad Z_n = \sqrt{n} \frac{\hat{\theta} - \theta}{b} \xrightarrow[n \rightarrow \infty]{D} Z \sim N(0, 1)$$

where  $b > 0$  is some (usually unknown) constant. Basically this is all we need. For a given “large”  $n$ , we interpret (0-1) as

$$(0-2) \quad \hat{\theta} \overset{\text{approximately}}{\sim} N\left(\theta, \frac{b^2}{n}\right)$$

where the expression  $\frac{b^2}{n}$  is often called the “asymptotic variance” of  $\hat{\theta}$  (the precise sense of which is given in (0-1)), and the square root is called the “(asymptotic) standard error”, SE,

$$(0-3) \quad SE(\hat{\theta}) = \frac{b}{\sqrt{n}}$$

Now, we cannot use (0-1) or (0-2) directly for inference when  $b$  is unknown. The only thing we need in order to deal with this problem, however, is a consistent estimator for  $b$ . So let  $\hat{b}$  be a consistent estimator for  $b$ . We then get from Slutsky's lemma (explain how) that

$$(0-4) \quad U_n = \sqrt{n} \frac{\hat{\theta} - \theta}{\hat{b}} \xrightarrow[n \rightarrow \infty]{D} Z \sim N(0, 1)$$

which we, for a given "large"  $n$ , may interpret as

$$(0-5) \quad \hat{\theta} \overset{\text{approximately}}{\sim} N\left(\theta, \frac{\hat{b}^2}{n}\right)$$

Note that the  $\hat{b}$  in the last expression can be understood as *the observed value* of  $\hat{b}$  (i.e., the estimate). In other words, for given "large"  $n$  we proceed as if the (asymptotic) standard error of  $\hat{\theta}$  is known and given by

$$(0-6) \quad SE(\hat{\theta}) = \frac{\hat{b}}{\sqrt{n}} \quad (\text{where } \hat{b} \text{ is the estimate and } \textit{not} \text{ the estimator.})$$

(This is the way Stata and other packages usually use the concept "standard error".)

The basis for inference can now be stated as:

For given "large"  $n$

$$(0-7) \quad U_n = \sqrt{n} \frac{\hat{\theta} - \theta}{\hat{b}} = \frac{\hat{\theta} - \theta}{SE(\hat{\theta})} \overset{\text{approximately}}{\sim} N(0, 1)$$

where  $\hat{b}$  is a consistent estimate of  $b$ .

**Exercise 1.** Identify  $b$  and  $\hat{b}$  in the following three situations:

i)  $X \sim \text{binomial}(n, p), \quad \theta = p, \quad \hat{\theta} = \hat{p} = X/n$

ii)  $X \sim \text{poisson}(\lambda t), \quad n = t, \quad \theta = \lambda, \quad \hat{\theta} = \hat{\lambda} = X/t$

iii)  $X_1, X_2, \dots, X_n \sim iid$  with  $E(X_i) = \mu = \theta, \quad \text{var}(X_i) = \sigma^2, \quad \hat{\theta} = \hat{\mu} = \bar{X}$

From (0-7) we can now construct confidence intervals (CI) for  $\theta$ , test hypotheses about  $\theta$ , calculate p-values etc.:

**Approximate  $1 - \alpha$  confidence interval for  $\theta$ :**

An approximate  $1 - \alpha$  CI for  $\theta$  is obtained directly from  $U_n$  in (0-7):

Let  $z_p$  be the upper  $p$ -point (quantile) in  $N(0, 1)$ , i.e.,  $P(Z > z_p) = p$ . Then:

$$\begin{aligned} 1 - \alpha &\approx P(-z_{\alpha/2} \leq U_n \leq z_{\alpha/2}) = P(-z_{\alpha/2} \leq \sqrt{n} \frac{\hat{\theta} - \theta}{\hat{b}} \leq z_{\alpha/2}) = \\ &= P\left(\hat{\theta} - z_{\alpha/2} \frac{\hat{b}}{\sqrt{n}} \leq \theta \leq \hat{\theta} + z_{\alpha/2} \frac{\hat{b}}{\sqrt{n}}\right) = \\ &= P\left(\hat{\theta} - z_{\alpha/2} \text{SE}(\hat{\theta}) \leq \theta \leq \hat{\theta} + z_{\alpha/2} \text{SE}(\hat{\theta})\right) \end{aligned}$$

Hence an approximate  $1 - \alpha$  CI for  $\theta$  is

$$(0-8) \quad \hat{\theta} \pm z_{\alpha/2} \text{SE}(\hat{\theta}) \quad \text{or} \quad \hat{\theta} \pm z_{\alpha/2} \frac{\hat{b}}{\sqrt{n}}$$

**Approximate  $\alpha$ -level tests for  $\theta$ :**

Consider the three following test problems in terms of null- and alternative hypotheses:

Problem	$H_0$	$H_1$
1	$\theta \leq \theta_0$	$\theta > \theta_0$
2	$\theta \geq \theta_0$	$\theta < \theta_0$
3	$\theta = \theta_0$	$\theta \neq \theta_0$

where  $\theta_0$  is a *known* hypothetical value. We can use the same test statistic for all three problems

$$(0-9) \quad W_n = \sqrt{n} \frac{\hat{\theta} - \theta_0}{\hat{b}} = \frac{\hat{\theta} - \theta_0}{\text{SE}(\hat{\theta})}$$

Notice the difference between  $U_n$  and  $W_n$  :

- $U_n$  is a non observable rv (since  $\theta$  is unknown) that has the same distribution ( $\approx N(0, 1)$ ) no matter if  $H_0$  is true or false.
- $W_n$  is an observable (its value can be calculated from the data) and its distribution is only  $\approx N(0, 1)$  if  $\theta$  should happen to be exactly equal to  $\theta_0$ .

The relationship between  $U_n$  and  $W_n$  is given by

$$W_n = \sqrt{n} \frac{\hat{\theta} - \theta_0}{\hat{b}} = \sqrt{n} \frac{\hat{\theta} - \theta + \theta - \theta_0}{\hat{b}} = U_n + \sqrt{n} \frac{\theta - \theta_0}{\hat{b}}$$

Hence, if  $\theta > \theta_0$ ,  $W_n$  behaves like a  $N(0, 1)$  variable plus something positive, if  $\theta < \theta_0$ ,  $W_n$  behaves like a  $N(0, 1)$  variable plus something negative, and if  $\theta = \theta_0$ ,  $W_n$  behaves just like a  $N(0, 1)$  variable.

The approximate test for problem 1 is therefore: “Reject  $H_0$  if  $W_n \geq c$ ”, where the critical value,  $c$ , is determined by the equation  $P_{\theta=\theta_0}(W_n \geq c) = \alpha$ . Since  $\theta = \theta_0$  implies that  $W_n$  is approximately  $N(0, 1)$ , we obtain  $c \approx z_\alpha$ . If  $w_{obs}$  denotes the observed value of  $W_n$  from the data, we get the (approximate) p-value:  $P_{\theta=\theta_0}(W_n \geq w_{obs}) \approx P(Z \geq w_{obs}) = 1 - \Phi(w_{obs})$ , where  $\Phi(x)$  is the cdf  $Z \sim N(0, 1)$ . Similarly for the other problems summarized in table 1:

**Table 1** Approximate  $\alpha$ -level tests

Problem	$H_0$	$H_1$	Reject $H_0$ when	Approximate P-value
1	$\theta \leq \theta_0$	$\theta > \theta_0$	$W_n \geq z_\alpha$	$P_{\theta=\theta_0}(W_n \geq w_{obs}) \approx P(Z \geq w_{obs})$
2	$\theta \geq \theta_0$	$\theta < \theta_0$	$W_n \leq -z_\alpha$	$P_{\theta=\theta_0}(W_n \leq w_{obs}) \approx P_{\theta=\theta_0}(Z \leq w_{obs})$
3	$\theta = \theta_0$	$\theta \neq \theta_0$	$ W_n  \geq z_{\alpha/2}$	$P_{\theta=\theta_0}( W_n  \geq  w_{obs} ) \approx P( Z  \geq  w_{obs} ) = 2P(Z \geq  w_{obs} )$

Note that we can achieve all this based on only the estimate and standard error obtained from e.g., a computer output, and a calculator, as illustrated in the following example:

**Example 0-1**

Suppose that we know from some computer output the estimate (based on say 40 observations) of a certain parameter  $\theta$  is  $\hat{\theta} = 1.598$  with standard error is  $SE = 0.356$ . In addition we know that the theory in (0-7) apply.

Then, for example, we may calculate an approximate 95% CI for  $\theta$  by

$$\hat{\theta} \pm 1.96 \cdot SE = 1.598 \pm 0.698 = [0.900, 2.296]$$

or a 90% CI

$$\hat{\theta} \pm 1.645 \cdot SE = 1.598 \pm 0.586 = [1.012, 2.184]$$

Suppose we want to test  $H_0 : \theta \leq 1$  against  $H_1 : \theta > 1$ . Then  $\theta_0 = 1$  above, and the test statistic is

$$W_n = \frac{\hat{\theta} - 1}{SE}$$

which gives the observed value

$$w_{obs} = \frac{1.598 - 1}{0.356} = 1.680$$

The p-value for this is approximately  $P_{\theta=1}(W_n > w_{obs}) \approx 1 - \Phi(1.680) = 0.046$

Thus  $H_0$  should be rejected on the 5% level but not on 1% level.

Most computer-packages calculate the so called “t” as well, given by  $t = \frac{\hat{\theta}}{SE(\hat{\theta})}$  (equal to

4.489 in this case). Notice that  $t$  is the same as the test statistic  $W_n$  when  $\theta_0 = 0$  in the problems above. The computer package also usually calculates the p-value for the two-sided hypothesis,  $H_0 : \theta = 0$  vs  $H_1 : \theta \neq 0$ , (under the somewhat cryptic heading like “p > |t|”), which corresponds to the p-value in the lower right cell of table 1. In this case the p-value in the output would be given as 0.000, which means that the p-value is less than 0.001. Thus  $\hat{\theta}$  is significantly (and highly so) different from 0 in this case.

*(End of example)*

In the (most common) case where  $\theta$  is one of several unknown parameters in the model, exactly the same principles of inference as above for  $\theta$  apply (in the case of asymptotic normality). The only difference is some technicalities in connection with the determination of standard errors of estimators from the asymptotic covariance matrix of the estimators, which will be the main topic (see section 3) in these lecture notes. As a background for this, we need a little bit of information (not in Rice) on random vectors and matrices, and on the multivariate normal distribution:

## 1 Random matrices

Let  $Y_{ij}$ ,  $i = 1, 2, \dots, m$ ,  $j = 1, 2, \dots, n$  be random variables (r.v.'s). The matrix

$$Y = \begin{pmatrix} Y_{11} & Y_{12} & \dots & Y_{1n} \\ Y_{21} & Y_{22} & \dots & Y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ Y_{m1} & Y_{m2} & \dots & Y_{mn} \end{pmatrix}$$

is called a random matrix ( with a joint  $mn$ -dimensional distribution,  $f(y_{11}, y_{12}, \dots, y_{mn})$  ).

The expected value of  $Y$  is *defined* as

$$(1) \quad E(Y) \stackrel{\text{def}}{=} \begin{pmatrix} E(Y_{11}) & E(Y_{12}) & \dots & E(Y_{1n}) \\ E(Y_{21}) & E(Y_{22}) & \dots & E(Y_{2n}) \\ \vdots & \vdots & \ddots & \vdots \\ E(Y_{m1}) & E(Y_{m2}) & \dots & E(Y_{mn}) \end{pmatrix}$$

The expectation satisfies the following rules (which follows directly from the definition (1) combined with the corresponding linear properties for the expectation in the scalar case):

- i.  $E(AY + C) = A \cdot E(Y) + C$   
where  $A, C$ , are any matrices of constants with dimensions compatible with  $Y$  (i.e.  $A \sim k \times m$ , and  $C \sim k \times n$ , where  $k$  is arbitrary).
- ii.  $E(AYB + C) = A \cdot E(Y) \cdot B + C$   
where  $A, B, C$  are any constant matrices compatible with  $Y$  in dimension so that the product and sum is well defined..
- iii.  $E(Y') = [E(Y)]'$  where  $A'$  denotes the transposed matrix

If  $Y = \begin{pmatrix} Y_1 \\ \vdots \\ Y_n \end{pmatrix}$  is a  $n$ -dimensional random vector, its expectation,  $\mu = E(Y)$  (sometimes written,  $\mu_Y$ ), is therefore the vector of individual expectations,

$$\mu = E(Y) = \begin{pmatrix} E(Y_1) \\ \vdots \\ E(Y_n) \end{pmatrix} = \begin{pmatrix} \mu_1 \\ \vdots \\ \mu_n \end{pmatrix}$$

Let  $\sigma_{ij} = E[(Y_i - \mu_i)(Y_j - \mu_j)] = \sigma_{ji}$  be the covariance between  $Y_i$  and  $Y_j$ . In particular we have  $\sigma_{ii} = E[(Y_i - \mu_i)^2] = \text{var}(Y_i)$ . The covariance matrix of  $Y$  (denoted as  $\text{cov}(Y)$ ) is defined as the matrix

$$\Sigma = \begin{pmatrix} \sigma_{11} & \cdots & \sigma_{1n} \\ \vdots & \ddots & \vdots \\ \sigma_{n1} & \cdots & \sigma_{nn} \end{pmatrix} = \begin{pmatrix} \text{var}(Y_1) & \cdots & \text{cov}(Y_1, Y_n) \\ \vdots & \ddots & \vdots \\ \text{cov}(Y_n, Y_1) & \cdots & \text{var}(Y_n) \end{pmatrix}$$

which can be expressed as

$$\begin{aligned} \text{cov}(Y) &= E[(Y - \mu)(Y - \mu)'] = E \begin{pmatrix} Y_1 - \mu_1 \\ \vdots \\ Y_n - \mu_n \end{pmatrix} (Y_1 - \mu_1, \dots, Y_n - \mu_n) = \\ &= E \begin{pmatrix} (Y_1 - \mu_1)^2 & \cdots & (Y_1 - \mu_1)(Y_n - \mu_n) \\ \vdots & \ddots & \vdots \\ (Y_n - \mu_n)(Y_1 - \mu_1) & \cdots & (Y_n - \mu_n)^2 \end{pmatrix} \stackrel{(1)}{=} \begin{pmatrix} \sigma_{11} & \cdots & \sigma_{1n} \\ \vdots & \ddots & \vdots \\ \sigma_{n1} & \cdots & \sigma_{nn} \end{pmatrix} = \Sigma \end{aligned}$$

### Example 1

Suppose that  $Y_1, Y_2, \dots, Y_n$  are *iid* with expectation  $E(Y_i) = \eta$  and  $\text{var}(Y_i) = \sigma^2$ . Then the vector  $Y' = (Y_1, \dots, Y_n)$  has expectation

$$E(Y) = \begin{pmatrix} \eta \\ \vdots \\ \eta \end{pmatrix} = \eta \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}$$

and covariance matrix (since  $\sigma_{ij} = \text{cov}(Y_i, Y_j) = 0$  for  $i \neq j$ ):

$$\text{cov}(Y) = \begin{pmatrix} \sigma^2 & 0 & \dots & 0 \\ 0 & \sigma^2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \sigma^2 \end{pmatrix} = \sigma^2 \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} = \sigma^2 I_n$$

where  $I_n$  is the  $n$ -dimensional identity matrix. (*End of example*)

If  $Y' = (Y_1, \dots, Y_n)$  is a random vector,  $A$  a  $p \times n$  constant matrix,  $b$  a constant  $p \times 1$  vector, we obtain from **i.-iii.** (and the fact that  $(BC)' = C'B'$  for matrices  $B$  and  $C$ ):

$$(2) \quad E(AY + b) = A \cdot E(Y) + b = A\mu + b$$

and

$$(3) \quad \text{cov}(AY + b) = A \cdot \text{cov}(Y) \cdot A' = A\Sigma A' \quad (\text{i.e. a } p \times p \text{ matrix})$$

which follows from

[remembering that for matrices  $A, B, C$  we always have  $A(B + C) = AB + AC$  and  $(B + C)A = BA + CA$  whenever the multiplication is well defined]

$$\begin{aligned} \text{cov}(AY + b) &= E[(AY + b - A\mu - b)(AY + b - A\mu - b)'] = E[(AY - A\mu)(AY - A\mu)'] = \\ &= E[A(Y - \mu)(Y - \mu)'A'] = A \cdot E[(Y - \mu)(Y - \mu)']A' = A\Sigma A' \end{aligned}$$

In particular, if  $Z$  is a linear combination of  $Y_1, \dots, Y_n$ , i.e.  $Z = a_0 + a_1Y_1 + \dots + a_nY_n$ , then  $Z$  can be written,  $Z = a_0 + a'Y$  where  $a' = (a_1, \dots, a_n)$  and  $Y = (Y_1, \dots, Y_n)'$ . Then (3) gives

$$(4) \quad \text{var}(Z) = \text{var}(a'Y) = a'\Sigma a \quad \text{where } \Sigma = \text{cov}(Y).$$

[**Proof:** Since  $Z = (a_1, \dots, a_n) \begin{pmatrix} Y_1 \\ \vdots \\ Y_n \end{pmatrix} = a'Y$  can be considered a  $1 \times 1$  matrix, we must have

that  $Z' = Z$ , and, therefore,  $\text{cov}(Z) = \text{var}(Z)$  (i.e.,

$$\text{cov}(Z) = E[(Z - E(Z))(Z - E(Z))'] = E[(Z - E(Z))^2] = \text{var}(Z)).$$

We then see that (4) is a special case of (3) with  $A = a'$  ]

### Example 2 Ordinary least squares (OLS)

To get an idea of the power of matrix notation, consider the standard multiple regression model with one response,  $Y$ , and  $p$  explanatory variables

$$(5) \quad Y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + u_i \quad \text{for } i = 1, 2, \dots, n$$

where, for simplicity, all  $x_{ij}$  are considered fixed, non random quantities, and the errors,  $u_1, u_2, \dots, u_n$  are assumed to be *iid* and normally distributed with expectation,  $E(u_i) = 0$  and  $\text{var}(u_i) = \sigma^2$ . We can write (5) in matrix form as follows

$$Y = \begin{pmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_n \end{pmatrix} = \begin{pmatrix} \beta_0 + \beta_1 x_{11} + \beta_2 x_{12} + \cdots + \beta_p x_{1p} \\ \beta_0 + \beta_1 x_{21} + \beta_2 x_{22} + \cdots + \beta_p x_{2p} \\ \vdots \\ \beta_0 + \beta_1 x_{n1} + \beta_2 x_{n2} + \cdots + \beta_p x_{np} \end{pmatrix} + \begin{pmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{pmatrix} = \begin{pmatrix} 1 & x_{11} & \cdots & x_{1p} \\ 1 & x_{21} & \cdots & x_{2p} \\ \vdots & \vdots & & \vdots \\ 1 & x_{n1} & \cdots & x_{np} \end{pmatrix} \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_p \end{pmatrix} + \begin{pmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{pmatrix}$$

The three matrices on the right we denote by  $X$ ,  $\beta$ , and  $u$  respectively. The model can now be written as

$$(6) \quad Y = X\beta + u$$

where  $X$  is the  $n \times p$  (so called) design matrix,  $\beta$  the  $(p+1) \times 1$  vector of regression coefficients, and  $u$  the  $n \times 1$  vector of errors. Since

$$E(u) = \begin{pmatrix} E(u_1) \\ E(u_2) \\ \vdots \\ E(u_n) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = \underline{0}$$

(where  $\underline{0}$  denotes a vector of zeroes), we get from **i.** (noting that  $X\beta$  is non random)

$$(7) \quad E(Y) = X\beta + E(u) = X\beta$$

The covariance matrix for  $Y$  becomes, since  $Y - X\beta = u$ , and using example 1,

$$(8) \quad \Sigma_Y = \text{cov}(Y) = E[(Y - X\beta)(Y - X\beta)'] = E[uu'] = \text{cov}(u) = \sigma^2 I_n$$

The OLS estimator,  $\hat{\beta}$ , for  $\beta$  is (by definition) obtained by minimizing the sum of squares

$$Q = \sum_{i=1}^n (Y_i - \beta_0 - \beta_1 x_{i1} - \cdots - \beta_p x_{ip})^2$$

with respect to  $\beta$ . Differentiating  $Q$  with respect to all the  $\beta_j$ 's, and setting the derivatives equal to 0, leads to (check if you wish) the following system of equations<sup>1</sup> that the  $\hat{\beta}_j$ 's must satisfy

$$\begin{aligned} n\hat{\beta}_0 + (\sum_i x_{i1})\hat{\beta}_1 + \cdots + (\sum_i x_{ip})\hat{\beta}_p &= \sum_i Y_i \\ (\sum_i x_{i1})\hat{\beta}_0 + (\sum_i x_{i1}^2)\hat{\beta}_1 + \cdots + (\sum_i x_{i1}x_{ip})\hat{\beta}_p &= \sum_i x_{i1}Y_i \\ &\dots \\ (\sum_i x_{ip})\hat{\beta}_0 + (\sum_i x_{ip}x_{i1})\hat{\beta}_1 + \cdots + (\sum_i x_{ip}^2)\hat{\beta}_p &= \sum_i x_{ip}Y_i \end{aligned}$$

Noting that the coefficients of the left side are exactly the elements in the  $(p+1) \times (p+1)$  matrix  $X'X$ , and that the right side, written as a vector, simply is  $X'Y$ , we can write the system more compactly as

$$X'X\hat{\beta} = X'Y$$

Assuming that  $X'X$  is non singular (which can be shown to be the case if no single  $x$ -variable can be written exactly as a linear combination of the other  $x$ -variables<sup>2</sup>, which is expressed by saying that there is no *exact collinearity* between the explanatory variables), we obtain the solution (the OLS estimator)

$$(9) \quad \hat{\beta} = (X'X)^{-1}X'Y$$

It is now easy to prove that  $\hat{\beta}$  is unbiased since, from **i.** and (7)

$$(10) \quad E(\hat{\beta}) = E[(X'X)^{-1}X'Y] \stackrel{i.}{=} (X'X)^{-1}X'E(Y) \stackrel{(7)}{=} (X'X)^{-1}X'X\beta = I_p\beta = \beta$$

Writing  $C = (X'X)^{-1}X'$ , we have  $\hat{\beta} = CY$ , and obtain the covariance matrix from (3) and (8)

[and also using the rule that the transposed of an inverse square matrix is the inverse of the transposed,  $[A^{-1}]' = (A')^{-1}$ , which is seen by taking the transposed of the equation,  $A \cdot A^{-1} = I$ . Remember also the  $AI_n = A$  for any  $p \times n$ -matrix  $A$ ,

<sup>1</sup> Often called *the normal equations* in the literature

<sup>2</sup> For example, if  $x_{i5} = 3 + x_{i1} - 2x_{i2}$  should happen to be true for all  $i = 1, 2, \dots, n$  in the data, there is exact collinearity between the variables  $x_1, x_2$ , and  $x_5$ . This would imply that  $X'X$  is singular.

and that, if  $c$  is a scalar, then  $c$  as factor can be taken outside a matrix product,  $A \cdot (cB) = cAB$ . ]

$$\text{cov}(\hat{\beta}) = \text{cov}(CY) \stackrel{(3)}{=} C\Sigma_Y C' \stackrel{(8)}{=} C(\sigma^2 I_n)C' = \sigma^2 CC' = \sigma^2 (X'X)^{-1} X'X (X'X)^{-1}$$

Hence

$$(11) \quad \text{cov}(\hat{\beta}) = \sigma^2 (X'X)^{-1} \quad (\text{End of example.})$$

**Exercise 2.** (Recommended (!) for obtaining a little bit of matrix training.)

Suppose that we are in a situation with only one (non-random) explanatory variable,  $x$ , so that the model in (5) reduces to

$$Y_i = \beta_0 + \beta_1 x_i + u_i \quad \text{for } i = 1, 2, \dots, n$$

where  $u_1, u_2, \dots, u_n$  are *iid* with  $E(u_i) = 0$  and  $\text{var}(u_i) = \sigma^2$

(a) Show that the determinant,  $D$ , of  $X'X$  is given by

$$D = n \sum_i x_i^2 - \left( \sum_i x_i \right)^2 = n \sum_i (x_i - \bar{x})^2$$

(b) Find the inverse  $(X'X)^{-1}$  and conclude that

$$\text{var}(\hat{\beta}_0) = \frac{\frac{1}{n} \sum_i x_i^2}{\sum_i (x_i - \bar{x})^2} \sigma^2 \quad \text{and} \quad \text{var}(\hat{\beta}_1) = \frac{\sigma^2}{\sum_i (x_i - \bar{x})^2}$$

(c) Show also by evaluating (9) that

$$\hat{\beta}_1 = \frac{\sum_i (x_i - \bar{x}) Y_i}{\sum_i (x_i - \bar{x})^2} = \frac{S_{xY}}{S_x^2} \quad \text{where}$$

$$S_{xY} = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y}) \quad \text{and} \quad S_x^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

(Note that this coincides with the estimation formula used for the corresponding parameter in *no-seminar-week exercise* for week 39)

## 2 Multinormal distributions

We say that the vector  $X' = (X_1, \dots, X_n)$  is (multi)normally distributed with expectation  $\mu = E(X)$ , and covariance matrix,  $\Sigma = \text{cov}(X)$  (written shortly  $X \sim N(\mu, \Sigma)$ ), if the joint pdf is given by

$$(12) \quad f(x_1, \dots, x_n | \mu, \Sigma) = \frac{1}{(2\pi)^{n/2} \sqrt{\det(\Sigma)}} e^{-\frac{1}{2}(x-\mu)'\Sigma^{-1}(x-\mu)} \quad \text{where } x = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \text{ and } \det(\Sigma)$$

means the determinant of  $\Sigma$ .

This distribution has a lot of convenient mathematical properties (see e.g. Greene, *Econometric Analysis*, chapter 3, for a summary), but here we only need the following:

$(13) \quad \text{If } X \sim N(\mu, \Sigma) \text{ and } A \text{ is a } p \times n \text{ constant matrix } (p \leq n) \text{ and } b \text{ a } p \times 1 \text{ constant vector, then } Y = AX + b \sim N(E(Y), \text{cov}(Y)) \stackrel{(2),(3)}{=} N(A\mu + b, A\Sigma A')$
--

[For proof see e.g. Greene chapter 3.]

In particular, this shows that all marginal distributions are also normal. For example, the marginal distribution of  $X_1, X_2$  is normal since

$$\begin{pmatrix} X_1 \\ X_2 \end{pmatrix} = AX \text{ where } A = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \end{pmatrix} \text{ which gives (check!)} \\ (14) \quad \begin{pmatrix} X_1 \\ X_2 \end{pmatrix} \sim N\left(E\begin{pmatrix} X_1 \\ X_2 \end{pmatrix}, \text{cov}\begin{pmatrix} X_1 \\ X_2 \end{pmatrix}\right) = N\left(\begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}, \begin{pmatrix} \sigma_{11} & \sigma_{12} \\ \sigma_{21} & \sigma_{22} \end{pmatrix}\right),$$

i.e. a bivariate normal distribution

**Exercise 3.** Show that the pdf in (14) as defined in (12), reduces to the bivariate normal density as defined in Example F in Rice section 3.3 (both editions). **[Hint:** Introduce the correlation,  $\rho$ , between  $X_1$  and  $X_2$ ,  $\rho = \sigma_{12} / \sqrt{\sigma_{11}\sigma_{22}}$ , implying

$\sigma_{12}^2 = \sigma_{11}\sigma_{22}\rho$ , and the determinant,  $\det(\text{cov}\begin{pmatrix} X_1 \\ X_2 \end{pmatrix}) = \sigma_{11}\sigma_{22} - \sigma_{12}^2 = \sigma_{11}\sigma_{22}(1 - \rho^2)$   
etc. ]

### Example 3 (Continuation of example 2)

The error vector,  $u$ , in (6) has expectation  $\underline{0}$  and covariance,  $\Sigma_u = \text{cov}(u) = \sigma^2 I_n$ . We see from (12) that saying that  $u \sim N(\underline{0}, \sigma^2 I_n)$  is the same as saying that  $u_1, u_2, \dots, u_n$  are *iid* and normally distributed with expectation,  $E(u_i) = 0$  and  $\text{var}(u_i) = \sigma^2$ . In fact, we have the determinant

$$\det(\Sigma_u) = \det(\sigma^2 I_n) = \det \begin{pmatrix} \sigma^2 & 0 & \dots & 0 \\ 0 & \sigma^2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \sigma^2 \end{pmatrix} = \sigma^{2n}.$$

and the exponent in (12) reduces to

$$-\frac{1}{2}(u - E(u))' \Sigma_u^{-1} (u - E(u)) = -\frac{1}{2} u' (\sigma^2 I_n)^{-1} u = -\frac{1}{2} u' \left( \frac{1}{\sigma^2} I_n \right) u = -\frac{1}{2\sigma^2} u' u = -\frac{1}{2\sigma^2} \sum_i u_i^2$$

Substituting in (12), shows that the joint distribution (12) reduces to the product of  $n$  one-dimensional  $N(0, \sigma^2)$ -distributions as the *iid* statement would imply.

By (13), (7), and (8) we obtain that  $Y$  is normally distributed,

$Y \sim N(E(Y), \text{cov}(Y)) = N(X\beta, \sigma^2 I_n)$ , and, by (13) again, that  $\hat{\beta} = (X'X)^{-1} X'Y$  is normally distributed

$$\hat{\beta} \sim N(E(\hat{\beta}), \text{cov}(\hat{\beta})) = N(\beta, \sigma^2 (X'X)^{-1})$$

[**Note:** For completeness sake it is worth mentioning that an unbiased (and consistent) estimator for  $\sigma^2$  is given by

$$\hat{\sigma}^2 = \frac{1}{n-p-1} \hat{u}' \hat{u} = \frac{1}{n-p-1} \sum_{i=1}^n \hat{u}_i^2$$

where  $\hat{u} = Y - X\hat{\beta}$  is the vector of the so called *residuals*.  $\hat{u}$  is an (unbiased) predictor of the non-observable error vector  $u$ . This enables us to get consistent

(and unbiased in the case of non-random  $X$ ) estimators of the covariance matrix of the regression estimators,  $\hat{\beta}$ .]

(End of example.)

### 3 On the asymptotic distribution for mle estimators (the multi parameter case)

In this section we will only describe how to determine the asymptotic distribution for the mle estimator in case there are several unknown parameters in the model, without going into details of derivations and proofs. A good summary of the theory can be found in chapter 4 of Greene's book, *Econometric Analysis*. See also Rice at the end of section 8.5.2.

Suppose that  $X_1, X_2, \dots, X_n$  are iid with  $X_i \sim f(x_i | \theta)$  (pdf), where  $\theta' = (\theta_1, \theta_2, \dots, \theta_r)$  is a  $r$ -dimensional vector of unknown parameters. Then the joint pdf is  $\prod_{i=1}^n f(x_i | \theta)$  and the log likelihood is

$$l(\theta) = \sum_{i=1}^n \ln f(x_i | \theta)$$

The mle estimator,  $\hat{\theta}$ , solves  $r$  equations

$$\sum_{i=1}^n \frac{\partial}{\partial \theta_j} \ln f(x_i | \hat{\theta}) = 0, \quad j = 1, 2, \dots, r$$

In order to define the  $r \times r$  Fisher information matrix that is needed in the asymptotic distribution of  $\hat{\theta}$ , we introduce

$$m_{ij}(\theta) = -E \frac{\partial^2 \ln f(X_i | \theta)}{\partial \theta_i \partial \theta_j} \quad i, j = 1, 2, \dots, r$$

Then the Fisher information matrix for one observation<sup>3</sup> is defined as

---

<sup>3</sup> Or one *observation vector* if we observe several variables per unit in the sample.

$$I(\theta) = \begin{pmatrix} m_{11}(\theta) & \cdots & m_{1r}(\theta) \\ \vdots & \ddots & \vdots \\ m_{r1}(\theta) & \cdots & m_{rr}(\theta) \end{pmatrix}$$

Under regularity conditions similar to the one-parameter case (see Greene for details), we have that the mle satisfies

$$\sqrt{n}(\hat{\theta} - \theta) \xrightarrow[n \rightarrow \infty]{D} N(\underline{0}, I(\theta)^{-1})$$

The definition of convergence in distribution for random vectors is similar but slightly more technical than the definition for the one-dimensional case, and we skip the details here (see Greene for a precise definition). However, the interpretation of the result is the same as in the one-dimensional case, i.e., that for large  $n$ ,

$$\hat{\theta} \stackrel{\text{approximately}}{\sim} N\left(\theta, \frac{1}{n}I(\theta)^{-1}\right)$$

Hence we can say that  $\hat{\theta}$  is asymptotically unbiased with asymptotic covariance matrix,  $(1/n)I(\theta)^{-1}$ . This matrix is unknown since  $\theta$  is unknown, but can be consistently estimated by replacing  $\theta$  by  $\hat{\theta}$  (or any other consistent estimator of  $\theta$ )<sup>4</sup>. [That  $\hat{\theta}$  is consistent means simply that  $\hat{\theta}_j \xrightarrow[n \rightarrow \infty]{P} \theta_j$  for all  $j = 1, 2, \dots, r$ ]. A generalization of Slutski's lemma to the multivariate case (details omitted), now allows us to conclude that, for large  $n$  (where  $\hat{\theta}_{obs}$  denotes the observed value of  $\hat{\theta}$ )

$$(15) \quad \boxed{\hat{\theta} \stackrel{\text{approximately}}{\sim} N\left(\theta, \frac{1}{n}I(\hat{\theta}_{obs})^{-1}\right)}$$

which is the important result that you should know. Using (13) we also have

$$(16) \quad A\hat{\theta} \stackrel{\text{approximately}}{\sim} N\left(A\theta, \frac{1}{n}A \cdot I(\hat{\theta}_{obs})^{-1} A'\right)$$

for any constant,  $p \times r$  matrix  $A$ .

---

<sup>4</sup> The continuity property of plim is still valid in the multivariate case, and the elements in  $I(\theta)^{-1}$  are continuous, which follows from the continuity of the elements of  $I(\theta)$ , expressing the inverse by determinants. Hence, the consistency of  $\hat{\theta}$  implies the consistency of  $I(\hat{\theta})^{-1}$ .

From this we get the following: Let  $k_{ij}(\theta)$  denote element  $i,j$  in  $I(\theta)^{-1}$ . Then the estimated asymptotic variance of  $\hat{\theta}_j$  is the  $j$ -th element on the main diagonal in the estimated covariance matrix, i.e.  $k_{jj}(\hat{\theta}_{obs})/n$ .

[Follows from (16). In fact, let  $a' = (0, \dots, 1, \dots, 0)$  where the 1 is in position  $j$  and zeroes elsewhere. Then from (16)

$$\hat{\theta}_j = a' \hat{\theta} \stackrel{\text{approx.}}{\sim} N\left(a' \theta, \frac{1}{n} a' I(\hat{\theta}_{obs})^{-1} a\right) = N\left(\theta_j, \frac{k_{jj}(\hat{\theta}_{obs})}{n}\right) ]$$

Hence, we obtain an approximate  $1-\alpha$  CI for  $\theta_j$ :  $\hat{\theta}_j \pm z_{\alpha/2} \sqrt{k_{jj}(\hat{\theta}_{obs})}/\sqrt{n}$  where  $z_{\alpha/2}$  is the upper  $\alpha/2$ -point in  $N(0, 1)$ .

**Example 4.** Assume we want a CI for the transformed parameter,  $\eta = \theta_1 - \theta_2$ . This we obtain from (16): Let  $b' = (1, -1, 0, \dots, 0)$ . Then, by (16),

$$\hat{\eta} = \hat{\theta}_1 - \hat{\theta}_2 = b' \hat{\theta} \stackrel{\text{approx.}}{\sim} N\left(b' \theta, \frac{b' I(\hat{\theta}_{obs})^{-1} b}{n}\right) = N\left(\theta_1 - \theta_2, \frac{1}{n} (k_{11}(\hat{\theta}_{obs}) + k_{22}(\hat{\theta}_{obs}) - 2k_{12}(\hat{\theta}_{obs}))\right)$$

which leads to the approximate  $1-\alpha$  CI for  $\theta_1 - \theta_2$ :

$$\hat{\theta}_1 - \hat{\theta}_2 \pm z_{\alpha/2} \frac{1}{\sqrt{n}} \sqrt{k_{11}(\hat{\theta}_{obs}) + k_{22}(\hat{\theta}_{obs}) - 2k_{12}(\hat{\theta}_{obs})}$$

[Note that all covariance matrices are symmetric. Hence  $k_{12}(\hat{\theta}) = k_{21}(\hat{\theta})$ . ]

(End of example.)

**Example 5 (On example C in Rice section 8.5 – precipitation data)**

Let  $X_i$  be the amount of precipitation for rainstorm no.  $i$ ,  $i = 1, 2, \dots, n$  ( $n = 227$  observations).

Model:  $X_1, X_2, \dots, X_n$  are *iid* with  $X_i \sim \Gamma(\alpha, \lambda)$ . The joint distribution is

$$X_1, X_2, \dots, X_n \sim \prod_{i=1}^n f(x_i | \alpha, \lambda) = \frac{\lambda^{n\alpha}}{\Gamma(\alpha)^n} (x_1 x_2 \dots x_n)^{\alpha-1} e^{-\lambda \sum x_i}$$

The log likelihood is

$$(17) \quad l(\alpha, \lambda) = n\alpha \ln \lambda + (\alpha - 1) \sum_i \ln x_i - \lambda \sum_i x_i - n \ln \Gamma(\alpha)$$

The first derivatives of  $l$  are

$$\frac{\partial l}{\partial \alpha} = n \ln \lambda + \sum_i \ln x_i - n \frac{\partial}{\partial \alpha} \ln \Gamma(\alpha)$$

$$\frac{\partial l}{\partial \lambda} = n \frac{\alpha}{\lambda} - \sum_i x_i$$

Setting the derivatives equal to zero and solving with respect to  $\alpha$  and  $\lambda$ , gives the mle estimators  $\hat{\alpha}$  and  $\hat{\lambda}$ .

[**Note.** There are no explicit formulas for the solution, they must be found by numerical iterations. For example, Excel works well in this case by the Solver module: Choose two cells for the arguments  $\alpha$  and  $\lambda$ , with start values e.g. at the moment estimates, and then a third cell for the function (17)<sup>5</sup>. Then use Solver to maximize (17). This can also be done in STATA by the ml-command, but slightly more involved.]

Using his program, Rice obtained the mle estimates.

$$\hat{\alpha} = 0,441 \quad \text{and} \quad \hat{\lambda} = 1,96$$

We want approximate 90% CI's for  $\alpha$  and  $\lambda$  based on the asymptotic normal distribution of  $\hat{\alpha}$  and  $\hat{\lambda}$ . In order to calculate the asymptotic standard errors we need the so called di- and trigamma functions defined by:

$$\text{Digamma function:} \quad \psi(\alpha) = \frac{\partial}{\partial \alpha} \ln \Gamma(\alpha)$$

$$\text{Trigamma function:} \quad \psi'(\alpha) = \frac{\partial^2}{\partial \alpha^2} \ln \Gamma(\alpha)$$

Both functions can be calculated in STATA (under the names digamma and trigamma).

---

<sup>5</sup> In Excel you can calculate  $\ln(\Gamma(\alpha))$  by the function GAMMALN.

We need the Fisher information matrix:

$$\ln f(X_i | \alpha, \lambda) = \alpha \ln \lambda - \ln \Gamma(\alpha) + (\alpha - 1) \ln X_i - \lambda X_i$$

giving

$$\frac{\partial \ln f}{\partial \alpha} = \ln \lambda - \psi(\alpha) + \ln X_i \quad \text{and} \quad \frac{\partial \ln f}{\partial \lambda} = \frac{\alpha}{\lambda} - X_i$$

Hence

$$\frac{\partial^2 \ln f}{\partial \alpha^2} = -\psi'(\alpha) \quad (\text{trigamma})$$

$$\frac{\partial^2 \ln f}{\partial \alpha \partial \lambda} = \frac{\partial^2 \ln f}{\partial \lambda \partial \alpha} = \frac{1}{\lambda}$$

$$\frac{\partial^2 \ln f}{\partial \lambda^2} = -\frac{\alpha}{\lambda^2}$$

Hence the Fisher information matrix for one observation

$$I(\alpha, \lambda) = -E \begin{pmatrix} -\psi'(\alpha) & \frac{1}{\lambda} \\ \frac{1}{\lambda} & -\frac{\alpha}{\lambda^2} \end{pmatrix} = \begin{pmatrix} \psi'(\alpha) & -\frac{1}{\lambda} \\ -\frac{1}{\lambda} & \frac{\alpha}{\lambda^2} \end{pmatrix}$$

The inverse of a symmetric  $2 \times 2$  matrix is

$$\begin{pmatrix} a & c \\ c & b \end{pmatrix}^{-1} = \frac{1}{ab - c^2} \begin{pmatrix} b & -c \\ -c & a \end{pmatrix}$$

Hence

$$I(\alpha, \lambda)^{-1} = \frac{1}{\frac{\alpha \psi'(\alpha)}{\lambda^2} - \frac{1}{\lambda^2}} \cdot \begin{pmatrix} \frac{\alpha}{\lambda^2} & \frac{1}{\lambda} \\ \frac{1}{\lambda} & \psi'(\alpha) \end{pmatrix} = \frac{1}{\alpha \psi'(\alpha) - 1} \cdot \begin{pmatrix} \alpha & \lambda \\ \lambda & \lambda^2 \psi'(\alpha) \end{pmatrix}$$

We obtain an estimate of this by substituting the mle,  $\hat{\alpha} = 0,441$  and  $\hat{\lambda} = 1,96$ , for  $\alpha$  and  $\lambda$  (skipping the index *obs* on the estimates)

$$I(\hat{\alpha}, \hat{\lambda})^{-1} = \frac{1}{\hat{\alpha}\psi'(\hat{\alpha}) - 1} \begin{pmatrix} \hat{\alpha} & \hat{\lambda} \\ \hat{\lambda} & \hat{\lambda}^2\psi'(\hat{\alpha}) \end{pmatrix} = \begin{pmatrix} 0,25903 & 1,15123 \\ 1,15123 & 13,82770 \end{pmatrix}$$

Here we found  $\psi'(\hat{\alpha}) = 6,128169$  from STATA by the command:

```
di trigamma(0.441)
```

From the theory we have that  $\begin{pmatrix} \hat{\alpha} \\ \hat{\lambda} \end{pmatrix} \overset{\text{approx.}}{\sim} N\left(\begin{pmatrix} \alpha \\ \lambda \end{pmatrix}, C\right)$ , where the asymptotic covariance is

$$C = \frac{1}{n} I(\hat{\alpha}, \hat{\lambda})^{-1} = \begin{pmatrix} 0,0011411 & 0,0050715 \\ 0,0050715 & 0,0609150 \end{pmatrix}$$

Hence the asymptotic standard errors

$$\text{se}(\hat{\alpha}) = \sqrt{0,0011411} = 0,03378 \quad \text{and} \quad \text{se}(\hat{\lambda}) = \sqrt{0,0609150} = 0,24681$$

According to the theory we then obtain approximate 90% CI for  $\alpha$  and  $\lambda$

$$\hat{\alpha} \pm 1,64 \cdot \text{se}(\hat{\alpha}) = 0,441 \pm (1,64)(0,03378) = [0,386, 0,496]$$

$$\hat{\lambda} \pm 1,64 \cdot \text{se}(\hat{\lambda}) = 1,96 \pm (1,64)(0,247) = [1,55, 2,37]$$

Rice (example E, section 8.5.3)) obtains approximate 90% CI's by the parametric bootstrap method:

$$\alpha: \quad [0,404, 0,523]$$

$$\lambda: \quad [1,46, 2,32]$$

The difference between the asymptotic intervals and the bootstrap intervals does not appear to be substantial. With as much as 227 observations it is to be expected that the asymptotic theory should work well.